

Clinical Appropriateness Guidelines

Molecular Testing of Solid and Hematologic Tumors and Malignancies

EFFECTIVE SEPTEMBER 6, 2021

ARCHIVED MARCH 6, 2022

This document has been archived because it has outdated information. It is for historical information only and should not be consulted for clinical use. Current versions of guidelines are available on the AIM Specialty Health website at <http://www.aimspecialtyhealth.com/>



8600 West Bryn Mawr Avenue
South Tower - Suite 800 Chicago, IL 60631
www.aimspecialtyhealth.com

Appropriate.Safe.Affordable
© 2021 AIM Specialty Health
GEN04-0921.1

Table of Contents

Scope	3
General Coverage Criteria	3
Conditions for which testing may be medically necessary.....	4
The following tests are not medically necessary	6
Specific Coverage Criteria	6
Breast Cancer Gene Expression Classifiers	6
Cell-Free Testing.....	8
Minimal Residual Disease (MRD) Testing.....	9
Targeted Molecular Testing for NTRK Fusions.....	9
Prostate Cancer (symptomatic cancer screening).....	10
CPT Codes.....	10
Background	20
Myeloproliferative Disorders	20
Gene Expression Classifiers	21
<i>Breast Cancer Gene Expression Classifiers</i>	21
<i>Prostate cancer (Post-Diagnosis Gene Expression Classifiers)</i>	23
<i>Cancer of Unknown Primary/Occult Primary Tumors</i>	24
Cell-Free Tumor Testing.....	24
Minimal Residual Disease (MRD) Genetic Testing	25
Tumor Agnostic Testing	26
<i>Microsatellite Instability and Immune Checkpoint Inhibitors</i>	26
<i>NTRK Fusion Testing</i>	26
Lung Cancer	27
Cancer Screening.....	28
<i>Indeterminate Thyroid Nodules</i>	28
<i>Prostate Cancer Early Screening</i>	30
Professional Society Guidelines.....	31
Selected References.....	33
Revision History.....	38

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Scope

This document addresses molecular testing and gene expression profiling of solid and hematologic tumors and malignancies (including cell free tumor DNA/circulating tumor cells/liquid biopsy testing) for the purpose of diagnosis, selecting chemotherapeutic agents and predicting risk, prognosis or recurrence of cancer. All tests listed in these guidelines may not require prior authorization; please refer to the health plan. For gene expression classifiers and polygenic risk scores not addressed in this policy, please refer to the Clinical Appropriateness Guidelines for Genetic Testing for Hereditary Cancer Susceptibility. For genetic testing used to guide chemotherapy treatment decisions, please refer to the Clinical Appropriateness Guidelines for Pharmacogenomic Testing and Genetic Testing for Thrombotic Disorders. In addition, testing required by a plan's pharmaceutical policies may be adjudicated by that plan's pharmaceutical guidelines.

General Coverage Criteria

Somatic tumor testing is medically necessary when all of the following criteria are met: (Please see below for conditions with additional specific criteria):

- Identification of the specific genetic variant or gene expression profile has been demonstrated through prospective research in peer-reviewed literature to improve diagnosis, management, or clinical outcomes for the individual's tumor type and disease characteristics
- Sample type (e.g., formalin-fixed, paraffin embedded (FFPE), cell-free tumor DNA, circulating tumor cells, etc.) has been proven to have clinical utility based on prospective evidence in peer-reviewed literature
- Testing methodology* has been clinically validated and is the most accurate method unless technical limitations (e.g., poor sample quality) necessitate the need for alternate testing strategies

*The testing methodology may target DNA and/or RNA.

In addition to the above criteria, somatic multi-gene panels for hematology-oncology indications are medically necessary when all of the following are met (please see additional criteria below for cell-free testing):

- Sequential testing of individual genes or biomarkers is not practical (i.e., limited tissue available, urgent treatment decisions pending) and more than one target is indicated
- Identification of genes or biomarkers on the panel has been demonstrated to improve diagnosis, management, or clinical outcomes for the individual's tumor type and disease characteristics
- The panel is targeted and limited to genes that are associated with the specific tumor type, unless otherwise specified in tumor site-specific criteria below

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

FDA companion diagnostics using NGS based panels may be considered medically necessary for the approved indication/medication when all of the following are met (see *Table 1. for specific approvable scenarios*):

- a more targeted test using any methodology is not available
- the patient does not otherwise meet criteria for treatment
- the patient meets criteria per the FDA label

Conditions for which testing may be medically necessary

Table 1. Molecular biomarkers that are medically necessary when the above criteria are met (list may not be all inclusive) (see *criteria below for cell-free testing*):

Indication	Molecular Studies
Acute Lymphoblastic Leukemia (ALL) <i>*See below for MRD testing criteria</i>	Targeted multi-gene panels
Acute Myelogenous Leukemia (AML) <i>*See below for MRD testing criteria</i>	Targeted multi-gene panels
B-Cell Lymphoma	EZH2, MYD88
Brain/Central Nervous System Cancers	IDH1, IDH2, MGMT, 1p/19q, ATRX, TERT, H3F3A, HIST1H3B, BRAF, RELA, TP53
Breast Cancer <i>*See below for gene expression classifier criteria</i> <i>*See below for cell-free testing criteria</i>	PIK3CA
Cholangiocarcinoma	FGFR2 FoundationOne® CDx (<i>tissue, for consideration of pemigatinib</i>)
Chronic Lymphocytic Leukemia (CLL)	TP53, IGHV, BTK, PLCG2
Chronic Myeloid Leukemia (CML)	BCR-ABL Targeted multi-gene panels
Colorectal Cancer Metastatic/Stage IV	BRAF, KRAS, NRAS Praxis Extended RAS panel (<i>tissue, for consideration of panitumumab</i>)
Endometrial Cancer	POLE
Essential Thrombocythemia or Thrombocytosis Platelet count $\geq 450 \times 10^9/L$	JAK2, CALR, MPL

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Gastrointestinal Stromal Tumors (GIST)	KIT, PDGFRA, SDHB, SDHC, SDHD, NF1, BRAF
Indeterminate Thyroid Nodule Bethesda Category III (AUS/FLUS) or Bethesda Category IV (FN/SFN) <i>*FNA samples with evidence of Hurthle cell pathology are excluded from coverage.</i>	BRAF, RAS, RET/PTC, PAX8/PPARc or ThyGeNEXT®/ThyraMIR™ or ThyroSeq® v3.0
Indeterminate Thyroid Nodule Bethesda Category III (AUS/FLUS) only <i>*FNA samples with evidence of Hurthle cell pathology are excluded from coverage.</i>	Afirma ² Genomic Sequence Classifier (GSC)
Melanoma (Cutaneous) Metastatic Melanoma (Stage III or Stage IV)	BRAF, KIT
Melanoma (Uveal)	EIF1AX, SF3B1, BAP1, PRAME, GNAQ, GNA11 or DecisionDx - Uveal Melanoma
Multiple Myeloma	Chromosomal Microarray Analysis (CMA) when cytogenetic (karyotype) and/or FISH analysis is uninformative
Myelodysplastic Syndrome	Targeted multi-gene panels
Neuroblastoma	Chromosomal Microarray Analysis (CMA), MYCN, ALK
Non-Small Cell Lung Cancer (NSCLC) Metastatic (Stage IIIB and above) <i>*See below for cell-free testing criteria</i>	ALK, BRAF, EGFR, ROS1, ERBB2 (HER2), MET, RET, KRAS Oncomine Dx Target Test (<i>tissue, for consideration of dabrafenib/trametinib, crizotinib, gefitinib, pralsetinib</i>)
Non-Small Cell Lung Cancer (NSCLC) Resected Stage IB – IIIA	EGFR
Ovarian, Fallopian Tube, or Primary Peritoneal Cancer <i>*See below for cell-free testing criteria</i>	BRCA1, BRCA2 myChoice® CDx (<i>tissue, for consideration of niraparib and/or olaparib</i>)
Polycythemia Vera Indication Includes ONE of the following (WHO criteria 2016): <ul style="list-style-type: none"> • Hemoglobin >16.5g/dL in men, >16.0g/dL in women • Hematocrit >49% in men, >48% in women • Increased red cell mass (RCM) 	JAK2, CALR, MPL
Primary Myelofibrosis Pre-PMF or suspicion for PMF based on 2016 WHO criteria	JAK2, CALR, MPL Targeted multi-gene panels (<i>when performed on bone marrow</i>)
Prostate Cancer (Suspected) Symptomatic Cancer Screening <i>*See criteria below.</i>	ConfirmMDx® ExoDx PCA3 SelectMDx

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Prostate Cancer Metastatic Castration-Resistant Prostate Cancer <i>*See below for cell-free testing criteria</i>	Targeted multi-gene panels FoundationOne® CDx (<i>tissue, for consideration of Olaparib</i>)
Tumor Agnostic/All Applicable Solid Tumors <i>*See NTRK criteria below</i>	Microsatellite Instability (MSI) NTRK 1/2/3 FoundationOne® CDx (<i>tissue, for consideration of pembrolizumab</i>)
T-Cell Lymphoma (Peripheral)	TET2, IDH1/IDH2, RHOA, DNMT3A, STAT3, STAT5B, TCR
Thyroid Cancer	BRAF, RET fusions

The following tests are not medically necessary

(list may not be all inclusive)

- Topographic genotyping (e.g., PancraGEN®)
- Whole exome tumor sequencing for any indication
- Whole genome tumor sequencing for any indication

In addition, testing of a genetic variant or profile correlated with a known therapy which does not have clinical utility for the specific tumor type and disease characteristics is not medically necessary.

Specific Coverage Criteria

Breast Cancer Gene Expression Classifiers

Breast cancer assays not listed below are considered not medically necessary.

Oncotype DX® Breast Recurrence Score Test is medically necessary to assess the need for adjuvant chemotherapy in the following individuals:

- Pre-menopausal women who are axillary-node negative or any axillary-node micrometastasis is no greater than 2.0 millimeters
- Post-menopausal women who are axillary-node negative or have no more than 3 positive lymph nodes
- Men who are axillary-node negative or have no more than 3 positive lymph nodes

AND all of the following criteria are met:

- Patient has undergone surgery and full pathological staging

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- Breast tumor is anatomic stage 1 or stage 2
- Histologic type is ductal, lobular, mixed (ductal/lobular), or metaplastic
- Tumor size >0.5 cm to ≤1.0 cm plus unfavorable histological features, defined as Nottingham grade 2-3 OR nuclear grade 3, or lymphovascular invasion **OR** tumor size 1.1-5.0 cm, any grade
- There is no evidence of distant metastatic breast cancer
- Breast tumor is estrogen and/or progesterone receptor-positive
- Breast tumor is HER2 receptor-negative
- Patient is a candidate for chemotherapy (i.e, chemotherapy not precluded due to other factors)
- Adjuvant chemotherapy is being considered and this testing is being ordered to assess recurrence risk to guide decision making as to whether or not adjuvant chemotherapy will be utilized
- No other breast gene expression classifier (GEC) has been performed

Prosigna™ PAM50, EndoPredict®, or Breast Cancer Index testing is medically necessary to assess the risk for recurrence in an individual when all of the following criteria are met:

- Patient has undergone surgery and full pathological staging
- Breast tumor is anatomic stage 1 or stage 2
- Histologic type is ductal, lobular, mixed (ductal/lobular), or metaplastic
- Tumor size >0.5 cm to ≤1.0 cm and intermediate or high grade (Grade 2 or 3) **OR** tumor size 1.1-5.0 cm, any grade
- Axillary-node status is negative or any axillary-node micrometastasis is no greater than 2.0 millimeters
- There is no evidence of distant metastatic breast cancer
- Breast tumor is estrogen or progesterone receptor-positive
- Breast tumor is HER2 receptor-negative
- Female patient is postmenopausal
- Patient is a candidate for chemotherapy (i.e, chemotherapy not precluded due to other factors)
- Adjuvant chemotherapy is being considered and this testing is being ordered to assess recurrence risk to guide decision making as to whether or not adjuvant chemotherapy will be utilized

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- No other breast GEC has been performed

MammaPrint® is medically necessary to assess the risk for recurrence in an individual when all of the following criteria are met:

- Patient has undergone surgery and full pathological staging
- Breast tumor is anatomic stage 1 or stage 2
- Histologic type is ductal, lobular, mixed (ductal/lobular), or metaplastic
- Node negative **OR** 1-3 positive node breast cancer
- Breast tumor is estrogen receptor positive and/or progesterone receptor positive
- Breast tumor is HER2-negative
- Patient is at high clinical risk for recurrence based on the MINDACT categorization
- Patient is a candidate for chemotherapy (i.e., chemotherapy not precluded due to other factors)
- Adjuvant chemotherapy is being considered and this testing is being ordered to assess recurrence risk to guide decision making as to whether or not adjuvant chemotherapy will be utilized
- No other breast GEC has been performed

Breast GEC testing is not medically necessary to guide decision making for extended endocrine therapy.

Cell-Free Testing

Cell-free testing (e.g., cfDNA, ctDNA, liquid biopsy) is medically necessary in the following scenarios when more targeted testing is not available:

- Metastatic Castrate-Resistant Prostate Cancer (mCRPC)
 - FoundationOne® Liquid CDx is medically necessary in men with metastatic castrate resistant prostate cancer (mCRPC) when the patient meets criteria per the FDA label for treatments for which this test has been approved as a companion diagnostic
 - Coverage is excluded when the patient already meets criteria without the need for additional testing (e.g., patient meets criteria based on known genetic results)
- Ovarian, Fallopian Tube, or Primary Peritoneal Cancer
 - FoundationOne® Liquid CDx is medically necessary in women with ovarian, fallopian tube, or primary peritoneal cancer when the patient meets criteria per the FDA label for treatment(s) for which this test has been approved as a companion diagnostic

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- Coverage is excluded when tissue-based testing is available
- Coverage is excluded when the patient already meets criteria without the need for additional testing (e.g., patient meets criteria based on known genetic results)
- Advanced or Metastatic Breast Cancer
 - theascreen[®] PIK3CA testing is medically necessary for advanced or metastatic breast cancer, using liquid biopsy if tumor is unavailable, when the patient meets criteria per the FDA label for treatments for which this test has been approved as a companion diagnostic
- Metastatic Non-Small Cell Lung Cancer (NSCLC)
 - At time of initial diagnosis
 - Targeted cell-free testing (e.g., cobas EGFR Mutation Test v2, ctDx Lung[™], Guardant360[®] CDx) is medically necessary in patients with metastatic NSCLC when tissue-based testing cannot be performed, e.g., insufficient tissue
 - At time of progression on an EGFR tyrosine kinase inhibitor (TKI) therapy
 - Targeted cell-free testing (i.e., cobas EGFR Mutation Test v2) is medically necessary in patients with metastatic NSCLC (excluding those patients who have progressed on osimertinib)

Minimal Residual Disease (MRD) Testing

NGS immunosequencing for MRD clone identification is covered when the following criteria is met:

- There is a confirmed diagnosis of B-cell acute lymphoblastic leukemia which is Philadelphia chromosome (BCR-ABL) negative
- Testing is performed on bone marrow

NGS minimal residual disease (MRD) testing for Philadelphia chromosome (BCR-ABL) negative B-cell ALL is covered when all of the following criteria are met:

- Immunosequencing at the time of diagnosis identified at least one clone for MRD tracking
- Complete cytologic remission is achieved
- Testing is performed on bone marrow

Targeted testing with prospective evidence of clinical utility for the tumor type and disease characteristics is covered.

Targeted Molecular Testing for NTRK Fusions

Targeted molecular testing for NTRK1/2/3 fusions is covered for any of the following indications:

- In tumors where NTRK fusions have a frequency of ~10% or greater (e.g., infantile fibrosarcoma, cellular congenital mesoblastic nephroma, secretory breast cancer, mammary

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

secretory carcinoma of the salivary gland, spitzoid melanoma, metastatic papillary thyroid cancer, analog pediatric high-grade glioma, or GIST when no KIT/PDGFR α /RAS pathogenic or likely pathogenic (P/LP) variant is identified)

- In solid tumors with positive NTRK IHC results or IHC is not possible for biomarker confirmation

Prostate Cancer (symptomatic cancer screening)

ExoDx (0005U) or SelectMDx (81479) is medically necessary for men ≥ 50 years considering initial biopsy when there is concern for prostate cancer as evidenced by a PSA of 3.1-10.0 ng/mL and none of the following:

- Treatment for benign prostatic hyperplasia in the past six months
- Treatment using a medication which impacts serum PSA levels within the past six months

PCA3 (81313) or ConfirmMDx (81551) is medically necessary for men ≥ 50 years with prior negative biopsy when repeat biopsy is being considered due to a PSA of 3.1-10.0 ng/mL.

Assays not listed above are considered not medically necessary.

CPT Codes

The following codes are associated with the guidelines in this document. This list is not all inclusive. Medical plans may have additional coverage policies that supersede these guidelines.

Covered when medical necessity criteria are met:

- | | |
|-------|---|
| 81162 | BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) (eg, hereditary breast and ovarian cancer) gene analysis; full sequence analysis and full duplication/deletion analysis (ie, detection of large gene rearrangements) |
| 81168 | CCND1/IGH (t(11;14)) (eg, mantle cell lymphoma) translocation analysis, major breakpoint, qualitative and quantitative, if performed |
| 81170 | ABL1 (ABL proto-oncogene 1, non-receptor tyrosine kinase) (eg, acquired imatinib tyrosine kinase inhibitor resistance), gene analysis, variants in the kinase domain |
| 81175 | ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; full gene sequence |

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 81176 ASXL1 (additional sex combs like 1, transcriptional regulator) (eg, myelodysplastic syndrome, myeloproliferative neoplasms, chronic myelomonocytic leukemia), gene analysis; targeted sequence analysis (eg, exon 12)
- 81191 NTRK1 (neurotrophic receptor tyrosine kinase 1) (eg, solid tumors) translocation analysis
- 81192 NTRK2 (neurotrophic receptor tyrosine kinase 2) (eg, solid tumors) translocation analysis
- 81193 NTRK3 (neurotrophic receptor tyrosine kinase 3) (eg, solid tumors) translocation analysis
- 81194 NTRK (neurotrophic-tropomyosin receptor tyrosine kinase 1, 2, and 3) (eg, solid tumors) translocation analysis
- 81206 BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; major breakpoint, qualitative or quantitative
- 81207 BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; minor breakpoint, qualitative or quantitative
- 81208 BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis; other breakpoint, qualitative or quantitative
- 81210 BRAF (B-Raf proto-oncogene, serine/threonine kinase) (eg, colon cancer, melanoma), gene analysis, V600 variant(s)
- 81218 CEBPA (CCAAT/enhancer binding protein [C/EBP], alpha) (eg, acute myeloid leukemia), gene analysis, full gene sequence
- 81219 CALR (calreticulin) (eg, myeloproliferative disorders), gene analysis, common variants in exon 9
- 81233 BTK (Bruton's tyrosine kinase) (eg, chronic lymphocytic leukemia) gene analysis, common variants (eg, C481S, C481R, C481F)
- 81235 EGFR (epidermal growth factor receptor) (eg, non-small cell lung cancer) gene analysis, common variants (eg, exon 19 LREA deletion, L858R, T790M, G719A, G719S, L861Q)
- 81236 EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit) (eg, myelodysplastic syndrome, myeloproliferative neoplasms) gene analysis, full gene sequence

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 81237 EZH2 (enhancer of zeste 2 polycomb repressive complex 2 subunit) (eg, diffuse large B-cell lymphoma) gene analysis, common variant(s) (eg, codon 646)
- 81245 FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; internal tandem duplication (ITD) variants (ie, exons 14, 15)
- 81246 FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia), gene analysis; tyrosine kinase domain (TKD) variants (eg, D835, I836)
- 81261 IGH@ (Immunoglobulin heavy chain locus) (eg, leukemias and lymphomas, B-cell), gene rearrangement analysis to detect abnormal clonal population(s); amplified methodology (eg, polymerase chain reaction)
- 81262 IGH@ (Immunoglobulin heavy chain locus) (eg, leukemias and lymphomas, B-cell), gene rearrangement analysis to detect abnormal clonal population(s); direct probe methodology (eg, Southern blot)
- 81263 IGH@ (Immunoglobulin heavy chain locus) (eg, leukemia and lymphoma, B-cell), variable region somatic mutation analysis
- 81264 IGK@ (Immunoglobulin kappa light chain locus) (eg, leukemia and lymphoma, B-cell), gene rearrangement analysis, evaluation to detect abnormal clonal population(s)
- 81270 JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) gene analysis, p.Val617Phe (V617F) variant
- 81272 KIT (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, gastrointestinal stromal tumor [GIST], acute myeloid leukemia, melanoma), gene analysis, targeted sequence analysis (eg, exons 8, 11, 13, 17, 18)
- 81273 KIT (v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog) (eg, mastocytosis), gene analysis, D816 variant(s)
- 81275 KRAS (Kirsten rat sarcoma viral oncogene homolog) (eg, carcinoma) gene analysis; variants in exon 2 (eg, codons 12 and 13)
- 81276 KRAS (Kirsten rat sarcoma viral oncogene homolog) (eg, carcinoma) gene analysis; additional variant(s) (eg, codon 61, codon 146)
- 81277 Cytogenomic neoplasia (genome-wide) microarray analysis, interrogation of genomic regions for copy number and loss-of-heterozygosity variants for chromosomal abnormalities
- 81278 IGH@/BCL2 (t(14;18)) (eg, follicular lymphoma) translocation analysis, major breakpoint region (MBR) and minor cluster region (mcr) breakpoints, qualitative or quantitative

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 81279 JAK2 (Janus kinase 2) (eg, myeloproliferative disorder) targeted sequence analysis (eg, exons 12 and 13)
- 81287 MGMT (O-6-methylguanine-DNA methyltransferase) (eg, glioblastoma multiforme) promoter methylation analysis
- 81301 Microsatellite instability analysis (eg, hereditary non-polyposis colorectal cancer, Lynch syndrome) of markers for mismatch repair deficiency (eg, BAT25, BAT26), includes comparison of neoplastic and normal tissue, if performed
- 81305 MYD88 (myeloid differentiation primary response 88) (eg, Waldenstrom's macroglobulinemia, lymphoplasmacytic leukemia) gene analysis, p.Leu265Pro (L265P) variant
- 81309 PIK3CA (phosphatidylinositol-4, 5-biphosphate 3-kinase, catalytic subunit alpha) (eg, colorectal and breast cancer) gene analysis, targeted sequence analysis (eg, exons 7, 9, 20)
- 81310 NPM1 (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, exon 12 variants
- 81311 NRAS (neuroblastoma RAS viral [v-ras] oncogene homolog) (eg, colorectal carcinoma), gene analysis, variants in exon 2 (eg, codons 12 and 13) and exon 3 (eg, codon 61)
- 81313 PCA3/KLK3 (prostate cancer antigen 3 [non-protein coding]/kallikrein-related peptidase 3 [prostate specific antigen]) ratio (eg, prostate cancer)
- 81314 PDGFRA (platelet-derived growth factor receptor, alpha polypeptide) (eg, gastrointestinal stromal tumor [GIST]), gene analysis, targeted sequence analysis (eg, exons 12, 18)
- 81315 PML/RARalpha, (t(15;17)), (promyelocytic leukemia/retinoic acid receptor alpha) (eg, promyelocytic leukemia) translocation analysis; common breakpoints (eg, intron 3 and intron 6), qualitative or quantitative
- 81316 PML/RARalpha, (t(15;17)), (promyelocytic leukemia/retinoic acid receptor alpha) (eg, promyelocytic leukemia) translocation analysis; single breakpoint (eg, intron 3, intron 6 or exon 6), qualitative or quantitative
- 81320 PLCG2 (phospholipase C gamma 2) (eg, chronic lymphocytic leukemia) gene analysis, common variants (eg, R665W, S707F, L845F)
- 81338 MPL (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; common variants (eg, W515A, W515K, W515L, W515R)
- 81339 MPL (MPL proto-oncogene, thrombopoietin receptor) (eg, myeloproliferative disorder) gene analysis; sequence analysis, exon 10

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 81340 TRB@ (T cell antigen receptor, beta) (eg, leukemia and lymphoma), gene rearrangement analysis to detect abnormal clonal population(s); using amplification methodology (eg, polymerase chain reaction)
- 81341 TRB@ (T cell antigen receptor, beta) (eg, leukemia and lymphoma), gene rearrangement analysis to detect abnormal clonal population(s); using direct probe methodology (eg, Southern blot)
- 81342 TRG@ (T cell antigen receptor, gamma) (eg, leukemia and lymphoma), gene rearrangement analysis, evaluation to detect abnormal clonal population(s)
- 81345 TERT (telomerase reverse transcriptase) (eg, thyroid carcinoma, glioblastoma multiforme) gene analysis, targeted sequence analysis (eg, promoter region)
- 81347 SF3B1 (splicing factor [3b] subunit B1) (eg, myelodysplastic syndrome/acute myeloid leukemia) gene analysis, common variants (eg, A672T, E622D, L833F, R625C, R625L)
- 81348 SRSF2 (serine and arginine-rich splicing factor 2) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variants (eg, P95H, P95L)
- 81351 TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; full gene sequence
- 81352 TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; targeted sequence analysis (eg, 4 oncology)
- 81353 TP53 (tumor protein 53) (eg, Li-Fraumeni syndrome) gene analysis; known familial variant
- 81357 U2AF1 (U2 small nuclear RNA auxiliary factor 1) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variants (eg, S34F, S34Y, Q157R, Q157P)
- 81360 ZRSR2 (zinc finger CCCH-type, RNA binding motif and serine/arginine-rich 2) (eg, myelodysplastic syndrome, acute myeloid leukemia) gene analysis, common variant(s) (eg, E65fs, E122fs, R448fs)
- 81450 Targeted genomic sequence analysis panel, hematolymphoid neoplasm or disorder, DNA and RNA analysis when performed, 5-50 genes (eg, BRAF, CEBPA, DNMT3A, EZH2, FLT3, IDH1, IDH2, JAK2, KRAS, KIT, MLL, NRAS, NPM1, NOTCH1), interrogation for sequence variants, and copy number variants or rearrangements, or isoform expression or mRNA expression levels, if performed
- 81479 SelectMDx
- 81518 Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 11 genes (7 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue,

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

algorithms reported as percentage risk for metastatic recurrence and likelihood of benefit from extended endocrine therapy

- 81519 Oncology (breast), mRNA, gene expression profiling by real-time RT-PCR of 21 genes, utilizing formalin-fixed paraffin embedded tissue, algorithm reported as recurrence score
- 81520 Oncology (breast), mRNA gene expression profiling by hybrid capture of 58 genes (50 content and 8 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence risk score
- 81521 Oncology (breast), mRNA, microarray gene expression profiling of 70 content genes and 465 housekeeping genes, utilizing fresh frozen or formalin-fixed paraffin-embedded tissue, algorithm reported as index related to risk of distant metastasis
- 81522 Oncology (breast), mRNA, gene expression profiling by RT-PCR of 12 genes (8 content and 4 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk score
- 81528 Oncology (colorectal) screening, quantitative real-time target and signal amplification of 10 DNA markers (KRAS mutations, promoter methylation of NDRG4 and BMP3) and fecal hemoglobin, utilizing stool, algorithm reported as a positive or negative result
- 81546 Oncology (thyroid), mRNA, gene expression analysis of 10,196 genes, utilizing fine needle aspirate, algorithm reported as a categorical result (eg, benign or suspicious)
- 81551 Oncology (prostate), promoter methylation profiling by real-time PCR of 3 genes (GSTP1, APC, RASSF1), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a likelihood of prostate cancer detection on repeat biopsy
- 81552 Oncology (uveal melanoma), mRNA, gene expression profiling by real-time RT-PCR of 15 genes (12 content and 3 housekeeping), utilizing fine needle aspirate or formalin-fixed paraffin-embedded tissue, algorithm reported as risk of metastasis
- 0005U Oncology (prostate) gene expression profile by real-time RT-PCR of 3 genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score
- 0018U Oncology (thyroid), microRNA profiling by RT-PCR of 10 microRNA sequences, utilizing fine needle aspirate, algorithm reported as a positive or negative result for moderate to high risk of malignancy
- 0022U Targeted genomic sequence analysis panel, non-small cell lung neoplasia, DNA and RNA analysis, 23 genes, interrogation for sequence variants and rearrangements, reported as presence/absence of variants and associated therapy(ies) to consider
- 0023U Oncology (acute myelogenous leukemia), DNA, genotyping of internal tandem duplication, p.D835, p.I836, using mononuclear cells, reported as detection or nondetection of FLT3 mutation and indication for or against the use of midostaurin

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 0026U Oncology (thyroid), DNA and mRNA of 112 genes, next-generation sequencing, fine needle aspirate of thyroid nodule, algorithmic analysis reported as a categorical result ("Positive, high probability of malignancy" or "Negative, low probability of malignancy")
- 0037U Targeted genomic sequence analysis, solid organ neoplasm, DNA analysis of 324 genes, interrogation for sequence variants, gene copy number amplifications, gene rearrangements, microsatellite instability and tumor mutational burden
- 0040U BCR/ABL1 (t(9;22)) (eg, chronic myelogenous leukemia) translocation analysis, major breakpoint, quantitative
- 0046U FLT3 (fms-related tyrosine kinase 3) (eg, acute myeloid leukemia) internal tandem duplication (ITD) variants, quantitative
- 0049U NPM1 (nucleophosmin) (eg, acute myeloid leukemia) gene analysis, quantitative
- 0111U Oncology (colon cancer), targeted KRAS (codons 12, 13, and 61) and NRAS (codons 12, 13, and 61) gene analysis utilizing formalin-fixed paraffin-embedded tissue
- 0154U Oncology (urothelial cancer), RNA, analysis by real-time RT-PCR of the FGFR3 (fibroblast growth factor receptor 3) gene analysis (ie, p.R248C [c.742C>T], p.S249C [c.746C>G], p.G370C [c.1108G>T], p.Y373C [c.1118A>G], FGFR3-TACC3v1, and FGFR3-TACC3v3) utilizing formalin-fixed paraffin-embedded urothelial cancer tumor tissue, reported as FGFR gene alteration status
- 0155U Oncology (breast cancer), DNA, PIK3CA (phosphatidylinositol-4,5-bisphosphate 3-kinase, catalytic subunit alpha) (eg, breast cancer) gene analysis (ie, p.C420R, p.E542K, p.E545A, p.E545D [g.1635G>T only], p.E545G, p.E545K, p.Q546E, p.Q546R, p.H1047L, p.H1047R, p.H1047Y), utilizing formalin-fixed paraffin embedded breast tumor tissue, reported as PIK3CA gene mutation status
- 0172U Oncology (solid tumor as indicated by the label), somatic mutation analysis of BRCA1 (BRCA1, DNA repair associated), BRCA2 (BRCA2, DNA repair associated) and analysis of homologous recombination deficiency pathways, DNA, formalin-fixed paraffin-embedded tissue, algorithm quantifying tumor genomic instability score
- 0177U Oncology (breast cancer), DNA, PIK3CA (phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha) gene analysis of 11 gene variants utilizing plasma, reported as PIK3CA gene mutation status
- 0179U Oncology (non-small cell lung cancer), cell-free DNA, targeted sequence analysis of 23 genes (single nucleotide variations, insertions and deletions, fusions without prior knowledge of partner/breakpoint, copy number variations), with report of significant mutation(s)
- 0239U Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free DNA, analysis of 311 or more genes, interrogation for sequence variants, including substitutions, insertions, deletions, select rearrangements, and copy number variations

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 0242U Targeted genomic sequence analysis panel, solid organ neoplasm, cell-free circulating DNA analysis of 55-74 genes, interrogation for sequence variants, gene copy number amplifications, and gene rearrangements
- 0245U Oncology (thyroid), mutation analysis of 10 genes and 37 RNA fusions and expression of 4 mRNA markers using next-generation sequencing, fine needle aspirate, report includes associated risk of malignancy expressed as a percentage
- ANY Clonoseq®

Codes that do not meet medical necessity criteria:

- 81327 SEPT9 (Septin9) (eg, colorectal cancer) promoter methylation analysis
- 81455 Targeted genomic sequence analysis panel, solid organ or hematolymphoid neoplasm, DNA and RNA analysis when performed, 51 or greater genes (eg, ALK, BRAF, CDKN2A, CEBPA, DNMT3A, EGFR, ERBB2, EZH2, FLT3, IDH1, IDH2, JAK2, KIT, KRAS, MLL, NPM1, NRAS, MET, NOTCH1, PDGFRA, PDGFRB, PGR, PIK3CA, PTEN, RET), interrogation for sequence variants and copy number variants or rearrangements, if performed
- 81504 Oncology (tissue of origin), microarray gene expression profiling of > 2,000 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as tissue similarity scores
- 81525 Oncology (colon), mRNA, gene expression profiling by real-time RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a recurrence score
- 81529 Oncology (cutaneous melanoma), mRNA, gene expression profiling by real-time RT-PCR of 31 genes (28 content and 3 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence risk, including likelihood of sentinel lymph node metastasis
- 81540 Oncology (tumor of unknown origin), mRNA, gene expression profiling by real-time RT-PCR of 92 genes (87 content and 5 housekeeping) to classify tumor into main cancer type and subtype, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported
- 81541 Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score
- 81542 Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 0011M Oncology, prostate cancer, mRNA expression assay of 12 genes (10 content and 2 housekeeping), RT-PCR test utilizing blood plasma and urine, algorithms to predict high-grade prostate cancer risk
- 0016M Oncology (bladder), mRNA, microarray gene expression profiling of 209 genes, utilizing formalin fixed paraffin-embedded tissue, algorithm reported as molecular subtype (luminal, luminal infiltrated, basal, basal claudin-low, neuroendocrine-like)
- 0017M Oncology (diffuse large B-cell lymphoma [DLBCL]), mRNA, gene expression profiling by fluorescent probe hybridization of 20 genes, formalin-fixed paraffin embedded tissue, algorithm reported as cell of origin
- 0045U Oncology (breast ductal carcinoma in situ), mRNA, gene expression profiling by realtime RT-PCR of 12 genes (7 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as recurrence score
- 0047U Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score
- 0048U Oncology (solid organ neoplasia), DNA, targeted sequencing of protein-coding exons of 468 cancer-associated genes, including interrogation for somatic mutations and microsatellite instability, matched with normal specimens, utilizing formalin-fixed paraffin-embedded tumor tissue, report of clinically significant mutation(s)
- 0050U Targeted genomic sequence analysis panel, acute myelogenous leukemia, DNA analysis, 194 genes, interrogation for sequence variants, copy number variants or rearrangements
- 0056U Hematology (acute myelogenous leukemia), DNA, whole genome nextgeneration sequencing to detect gene rearrangement(s), blood or bone marrow, report of specific gene rearrangement(s)
- 0057U Oncology (solid organ neoplasia), mRNA, gene expression profiling by massively parallel sequencing for analysis of 51 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a normalized percentile rank
- 0069U Oncology (colorectal), microRNA, RT-PCR expression profiling of miR-31-3p, formalin-fixed paraffin-embedded tissue, algorithm reported as an expression score
- 0089U Oncology (melanoma), gene expression profiling by RTqPCR, PRAME and LINC00518, superficial collection using adhesive patch(es)
- 0090U Oncology (cutaneous melanoma), mRNA gene expression profiling by RT-PCR of 23 genes (14 content and 9 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a categorical result (ie, benign, indeterminate, malignant)

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 0113U Oncology (prostate), measurement of PCA3 and TMPRSS2-ERG in urine and PSA in serum following prostatic massage, by RNA amplification and fluorescence-based detection, algorithm reported as risk score
- 0114U Gastroenterology (Barrett's esophagus), VIM and CCNA1 methylation analysis, esophageal cells, algorithm reported as likelihood for Barrett's esophagus
- 0120U Oncology (B-cell lymphoma classification), mRNA, gene expression profiling by fluorescent probe hybridization of 58 genes (45 content and 13 housekeeping genes), formalin-fixed paraffin-embedded tissue, algorithm reported as likelihood for primary mediastinal B-cell lymphoma (PMBCL) and diffuse large B-cell lymphoma (DLBCL) with cell of origin subtyping in the latter
- 0153U Oncology (breast), mRNA, gene expression profiling by next-generation sequencing of 101 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a triple negative breast cancer clinical subtype(s) with information on immune cell involvement
- 0171U Targeted genomic sequence analysis panel, acute myeloid leukemia, myelodysplastic syndrome, and myeloproliferative neoplasms, DNA analysis, 23 genes, interrogation for sequence variants, rearrangements and minimal residual disease, reported as presence/absence
- 0204U Oncology (thyroid), mRNA, gene expression analysis of 593 genes (including BRAF, RAS, RET, PAX8, and NTRK) for sequence variants and rearrangements, utilizing fine needle aspirate, reported as detected or not detected
- 0208U Oncology (medullary thyroid carcinoma), mRNA, gene expression analysis of 108 genes, utilizing fine needle aspirate, algorithm reported as positive or negative for medullary thyroid carcinoma
- 0211U Oncology (pan-tumor), DNA and RNA by next-generation sequencing, utilizing formalin-fixed paraffin-embedded tissue, interpretative report for single nucleotide variants, copy number alterations, tumor mutational burden, and microsatellite instability, with therapy association
- 0228U Oncology (prostate), multianalyte molecular profile by photometric detection of macromolecules adsorbed on nanosponge array slides with machine learning, utilizing first morning voided urine, algorithm reported as likelihood of prostate cancer
- 0229U BCAT1 (Branched chain amino acid transaminase 1) or IKZF1 (IKAROS family zinc finger 1) (eg, colorectal cancer) promoter methylation analysis
- 0235U PTEN (phosphatase and tensin homolog) (eg, Cowden syndrome, PTEN hamartoma tumor syndrome), full gene analysis, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 0238U Oncology (Lynch syndrome), genomic DNA sequence analysis of MLH1, MSH2, MSH6, PMS2, and EPCAM, including small sequence changes in exonic and intronic regions, deletions, duplications, mobile element insertions, and variants in non-uniquely mappable regions
- 0244U Oncology (solid organ), DNA, comprehensive genomic profiling, 257 genes, interrogation for single-nucleotide variants, insertions/deletions, copy number alterations, gene rearrangements, tumor-mutational burden and microsatellite instability, utilizing formalin-fixed paraffin embedded tumor tissue
- ANY Guardant360® LDT for any indication (Guardant Health, Inc.)

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright by the American Medical Association. All Rights Reserved. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Background

Somatic genetic testing for the purpose of cancer management guidance is a rapidly evolving field of molecular medicine. Genetic testing of a solid tumor or hematologic neoplasm can provide important information regarding the prognosis, risk for recurrence or help predict response to chemotherapeutic agents. In addition, genetic testing of tissue (e.g., blood) or stool, for evidence of a tumor, is becoming an important tool in the early detection of cancer. While this is an area of ongoing research, clinical validity and utility is proven for only a subset of companion diagnostic genetic tests at this time.

Myeloproliferative Disorders

Myeloproliferative disorders, or myeloproliferative neoplasms (MPNs), are a group of conditions that cause abnormal growth of blood cells in the bone marrow. They include polycythemia vera (PV), essential thrombocythemia or thrombocytosis, pre-primary myelofibrosis, primary myelofibrosis, chronic myelogenous leukemia, and chronic neutrophilic leukemia. The diagnosis of an MPN is suspected based upon clinical, laboratory, and pathological findings, including bone marrow morphology and certain pathogenic/likely pathogenic (P/LP) variants. Patients with MPNs may be clinically asymptomatic, but MPNs confer a risk for progression to acute myeloid leukemia, also called blast-phase MPN (Lasho et al. 2018).

JAK2, CALR, and MPL are genes involved in the growth and survival of various cell types. The presence of somatic driver mutations within these genes is part of the World Health Organization diagnostic criteria for MPNs, and molecular testing may be necessary to confirm a diagnosis. Chronic myelogenous leukemia (CML) is distinguished from the other MPNs by the presence of a BCR-ABL1 fusion gene. Targeted genetic testing of the JAK2, CALR, and MPL genes may be helpful in individuals who would not otherwise meet diagnostic criteria. There is some evidence that non-driver P/LP variants in additional genes such as ASXL1, TET2, and TP53 can help to predict a poor prognosis for some patients with MPNs, but the utility of testing these genes is not fully established (McClure et al. 2018; Grinfeld et al. 2018). Importantly, P/LP variants in many of these genes have also been detected in

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

older individuals with no other clinical evidence of myeloid disease, a scenario known as clonal hematopoiesis of indeterminate potential (CHIP). Therefore, genetic testing should only be performed when there is reasonable clinical suspicion of disease. At this time, variants in other genes associated with MPNs, including P/LP variants within ASXL1, TET2, SRSF2, U2AF1, IDH1/IDH2, TP53, DNMT3A, IKZF1, LNK, SF3B1, EZH2, CBL, and SETBP1, are recommended only in the evaluation for primary and pre-primary myelofibrosis.

MPNs are related to, but distinct from, myelodysplastic syndromes (MDS). In general, MDS are characterized by ineffective or dysfunctional blood cells with an increased risk of transformation to acute myeloid leukemia (AML), while MPNs are characterized by an increase in the number of blood cells. MDS typically first present as cytopenia(s) or dysplasia in one or more hematopoietic cell lines in the bone marrow. The development and transformation of MDS is driven by somatic variants in genes related to RNA splicing, epigenome regulation, myeloid transcriptional coordination, DNA damage and stress responses, and/or growth factor signaling. The diagnosis of an MDS relies on incorporating clinical features, peripheral blood and bone marrow findings, and cytogenetic analysis. Molecular testing may be appropriate in select scenarios when a diagnosis already exists and testing will help clarify prognostic category, which can help guide the treatment pathway.

Gene Expression Classifiers

Breast Cancer Gene Expression Classifiers

While NGS panels are not currently recommended for use to guide chemotherapeutic treatment decisions, molecular testing may be used to predict prognosis and recurrence risk for breast cancer. Along with a patient's age and comorbidities, the strongest prognostic factors to predict future recurrence or death from breast cancer include patient age, comorbidity, tumor size, tumor grade, number of involved axillary lymph nodes, and HER2 tumor status (Cao 2016).

Breast cancer gene expression profiling refers to testing performed on breast cancer tumor tissue to identify expression levels of sets of genes that, taken together, may predict recurrence risk and/or treatment response.

The American Society of Clinical Oncology (ASCO) published recommendations on the management of male breast cancer (2019) that revealed high-level consensus for similar management in men and women regarding the use of gene expression profile testing to guide adjuvant treatment decision making (e.g., Oncotype DX and prognostic tests). ASCO (2016) recommends use of the Oncotype Dx[®] assay to guide decisions on adjuvant chemotherapy in patients treated with tamoxifen who are node-negative, HER2 negative, and estrogen-receptor positive (Harris et al. 2016).

Sufficient data supports the use of the Oncotype Dx[®] assay for recurrence risk prediction and determination of adjuvant chemotherapy for:

- Early anatomic stage (I or II) invasive breast cancer, AND
- Axillary lymph node negative / no evidence of distant metastatic breast cancer / any axillary-node micrometastasis is 2 mm or less, AND
- Estrogen receptor positive AND
- HER2 receptor negative AND
- Patients who are candidates for adjuvant chemotherapy

In 2019 ASCO updated their guidelines again by incorporating data from the TAILORx trial. No changes were made to the criteria for whom should be offered OncotypeDX testing; however, adjuvant systemic treatment options were further delineated based on OncotypeDX recurrence score (Andre et al. 2019; Sparano et al. 2019).

The 2016 ASCO practice guideline published in the *Journal of Clinical Oncology* supports the use of certain tumor biomarker assays beyond the Oncotype Dx[®] Breast assay described above, in select populations to guide treatment. Importantly, these recommendations are based on review of evidence in which no true prospective trials have been performed (Harris et al. 2016). Specifically, ASCO supports the use of the following tests in the outlined scenarios:

- EndoPredict[®] for women with ER/PR-positive, HER2-negative, node-negative breast cancer to guide decisions on adjuvant systemic chemotherapy. This is an evidence-based recommendation with reported intermediate evidence quality, and a moderate strength of recommendation
- Prosigna[™] PAM50 Breast Cancer Prognostic Gene Signature Assay for women with ER/PR-positive, HER2-negative, node-negative breast cancer to be used in conjunction with other clinicopathologic variables to guide decisions on adjuvant systemic therapy. This is an evidence-based recommendation with reported high-quality evidence and a strong strength of recommendation
- Breast Cancer Index[®] (BCI) for women with ER/PR-positive, HER2-negative, node-negative breast cancer to guide decisions on adjuvant systemic therapy. This is an evidence-based recommendation with intermediate quality evidence, and a moderate strength of recommendation

ASCO published a special addendum (Krop et al. 2017) regarding use of MammaPrint[®] for women with hormone receptor- positive, HER2-negative, node negative and node positive tumors based on preliminary MINDACT data (Cardoso et al. 2016) that was reaffirmed in 2019 (Henry et al. 2019). The prior recommendation for this group [women with HR+, HER2- (node positive or node-negative) breast cancer] was that the clinician should not use MammaPrint[®] to guide decisions on adjuvant systemic chemotherapy. The 2017 updated guideline separates this group into 3 categories and recommendations:

- **Recommendation 1.1.1:** MammaPrint[®] assay may be used for women with hormone receptor- positive, HER2-negative, node negative cancer who are considered high clinical risk per MINDACT categorization to inform decision making regarding withholding adjuvant systemic chemotherapy due to its ability to identify a good prognosis population with potentially limited chemotherapy benefit. (Evidence Quality: High and Strength of Recommendation: Strong)
- **Recommendation 1.1.2:** MammaPrint[®] assay should not be used for women with hormone receptor- positive, HER2-negative, node negative cancer who were considered low clinical risk per MINDACT categorization because women in the low clinical risk category had excellent outcomes and did not seem to benefit from chemotherapy even with a genomically high risk cancer. (Evidence Quality: High and Strength of Recommendation: Strong)
- **Recommendation 1.2.1:** MammaPrint[®] assay may be used in patients with hormone receptor- positive, HER2-negative, node positive (with 1-3 positive nodes) cancer and at high clinical risk per MINDACT categorization to inform decision making regarding

withholding adjuvant systemic chemotherapy because of its ability to identify a good prognosis population with potentially limited chemotherapy benefit. Patients should be informed that the benefit of chemotherapy cannot be excluded, particularly in patients with more than one involved lymph node. (Evidence Quality: High; Strength of Recommendation: Moderate)

While the clinical utility of the OncotypeDx Recurrence Score (RS) has been established in node negative, HR positive, HER2-negative patients with breast cancer; results from the RxPonder trial have been needed to establish its utility in node positive patients with similar breast cancer characteristics. An independent safety monitoring committee recommended reporting findings publicly prior to the final analysis after noting a surprising and clear pattern of benefit for postmenopausal women (Kalinsky et al. 2020). A significant association between recurrence score and chemotherapy benefit was found with menopausal status ($p=0.004$). While patients will still be followed for 15 years, the current data suggest adjuvant therapy can be ET alone in postmenopausal patients with 1-3 positive nodes and a $RS \leq 25$. The opposite is true for premenopausal women after data revealed invasive disease-free survival (IDFS) benefit for chemoendocrine therapy (CET) (Kalinsky et al. 2020).

The following tests are not supported within the ASCO practice guideline under any circumstances at this time: MammoStrat® or any assays performed using circulating tumor cells or tumor-infiltrating lymphocytes.

Prostate cancer (Post-Diagnosis Gene Expression Classifiers)

The American Urological Association (AUA), ASTRO and the Society of Urologic Oncology (SUO) published guidelines in 2018 for risk stratification, shared decision making, and care options for clinically localized prostate cancer. It is notable that these guidelines do not include a recommendation for genomic testing of prostate tumor samples, and instead use Gleason score, PSA, and clinical stage in the risk stratification and assessment of treatment options. The authors state that no genomic tests have yet been validated as providing substantial benefit in the active surveillance population (Sanda et al. 2017; Sanda et al. 2018). The European Association of Urology recently created (and externally validated) a simple risk stratification system to help identify men at high risk for biochemical recurrence; this schema uses Gleason score and PSA levels - notably absent is the incorporation of any gene expression assays (Van den Broeck et al. 2020). The American Society of Clinical Oncologists (ASCO) recently released recommendations supporting the use of commercially available molecular biomarkers in situations in which the assay results, coupled with other routine clinical factors, would be likely to change medical management (Eggerer et al. 2019). However, the ASCO statement notes that “there is a paucity of prospective studies assessing short- and long-term outcomes of patients when these markers are integrated into clinical decision making.” (Eggerer et al. 2020).

Naryan et al. (2017), performed an evidence-based review for biomarker assays used for prostate cancer. The group reviewed Prolaris® and Oncotype DX® Prostate and commented that although these tests have been incorporated into NCCN Guidelines® and may be beneficial for men with low-volume Gleason 6 disease on biopsy, these tests have not been thoroughly studied in minority populations, and it is unclear how initial test results may change with repeat assessments. They recommend that these tests should be used with discretion as they add to the cost of prostate cancer care and that providers should discuss the indications and limitations thoroughly with their patients (Narayan et al. 2017). Similarly, Lamy et al. (2017) performed a systematic review of prostate cancer biomarkers and concluded the Prostate Health Index and the 4K score have the highest level of evidence in predicting which cancers may be more aggressive. They also note that other assays, including OncotypeDX®

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Prostate, Prolaris[®], and Decipher[®] Prostate, are promising but need further evidence to confirm their clinical validity.

For men with metastatic castrate-resistant prostate cancer (mCRPC), there has been interest in the use of testing of circulating tumor cells (CTCs) for a splice site variant in the androgen receptor gene, AR-V7, to help guide therapeutic intervention, particularly in the setting of progression on androgen receptor signaling inhibitors (ARSI) such as abiraterone or enzalutamide. This potential biomarker has been extensively studied, with conflicting results (Kretschmer et al. 2017; Scher et al. 2018; Armstrong et al. 2019; Abida et al. 2019). While there is prospective evidence demonstrating men affected by mCRPC with the AR-V7 variant in CTCs have worse outcomes when treated with enzalutamide/abiraterone, there is not currently prospective evidence they do better on an alternate therapy. More evidence is needed to show AR-V7 is a reliable biomarker to predict response to improved outcomes in this regard. ASCO guidelines indicate that there is no evidence of clinical utility and little evidence of clinical validity of ctDNA assays in early-stage cancer, treatment monitoring, or residual disease detection (Merker et al. 2018).

Cancer of Unknown Primary/Occult Primary Tumors

Occult primary tumors, or cancers of unknown primary, are defined as histologically proven metastatic malignant tumors whose primary site cannot be identified by a standard diagnostic workup during pretreatment evaluation. These may have a wide clinical presentation and typically a poor prognosis (Binder et al. 2018). It has been proposed that more intensive diagnostic studies aimed at identifying the primary cancer site is important to guide disease-oriented therapy. Several laboratories offer gene expression profiling (GEP) or NGS tests to aid in the identification of the tissue of origin of a metastatic tumor (Binder et al. 2018). The current literature evaluating molecular testing in the diagnosis and management of occult primaries has focused much more on establishing the tissue of origin rather than establishing whether such identification leads to better outcomes for patients. Although these results may have diagnostic benefit, there is limited evidence that management changes based on results impact patient survival. A randomized phase II trial found no improvement in 1-year survival between patients who were treated with site-specific therapies based on GEP results and patients who were treated with empirical chemotherapy (Hayashi et al. 2019).

Multiple professional societies have commented on the limited evidence of clinical utility for molecular testing to identify the origin of occult primary cancers. The European Society for Medical Oncology (ESMO) notes the potential promise of molecular assays to assist with tissue of origin identification for cancers of unknown primary; however, the ESMO clinical practice guideline goes on to note insufficient evidence related to further using assay-predicted tumor type to guide primary site-specific therapy (Fizazi et al. 2015).

Cell-Free Tumor Testing

Tumor testing for recommended markers is not always possible, primarily due to an inadequate tissue sample. It is estimated that 15% of patients with NSCLC who undergo biopsy have an inadequate sample for molecular testing (Douillard et al. 2014). In addition, many patients with late-stage metastatic cancer may be poor candidates for biopsy.

There has been growing interest and research into alternative blood-based methodologies for assessing tumor P/LP variant status, including cell-free plasma-based tests. An example is cell-free tumor DNA (ctDNA) testing which is commonly employed because ctDNA is easier to isolate and, with the increasing capabilities of next-generation sequencing, it offers an alternate opportunity to assess

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

somatic tumor-specific P/LP variants. While several studies have shown that ctDNA is not as sensitive or specific as direct tumor testing (Janku et al. 2016; Zhang et al. 2016), positive results are generally assumed to be accurate enough to use in treatment planning (Madison et al. 2020). There are potential applications where ctDNA testing might be indicated (e.g., when a biopsy sample is insufficient, when repeat biopsy is overly risky, or when chemotherapy response has changed and there is a concern for intra- or inter-tumor heterogeneity) to provide information about the molecular status of a tumor (Rolfo et al. 2018). It has also been proposed that ctDNA may improve minimal residual disease monitoring (Levy et al. 2016).

Cell-free tumor DNA analysis is still an active area of research and monitoring of performance data will be ongoing. Currently, there are select clinical scenarios with sufficient evidence to allow cell-free tumor DNA analysis to help guide therapeutic decision making: metastatic NSCLC, metastatic breast cancer, metastatic castrate-resistant prostate cancer or ovarian cancer. Utility has not yet been proven in other clinical scenarios including the use of methylation of the SEPT9 gene (mSEPT9) for colon cancer screening. Concerns remain regarding the poor specificity of this testing methodology for colon cancer, and the USPSTF along with the American Cancer Society do not recommend the use of SEPT9 for colorectal cancer screening in any scenarios (Rex et al. 2017; Wolf et al. 2018). Additional studies of circulating tumor DNA have not shown that this technique is able to reliably detect other colon tumor-related P/LP variants (Myint et al. 2018; Liebs et al. 2019). In general, there is also insufficient evidence to recommend coverage of plasma-based testing (ctDNA) over tumor-based testing when an appropriate tumor sample is available (Rolfo et al. 2020). ctDNA testing may be reasonable to aid treatment selection at the time of diagnosis or to guide treatment decision making when progression occurs.

Minimal Residual Disease (MRD) Genetic Testing

Minimal residual disease (MRD) refers to persistence of low levels of tumor cells after a patient appears to have achieved complete remission from drug therapy or HCT. MRD may also be referred to as "measurable residual disease." Detecting MRD after apparent treatment success is of great interest to define patients at risk for relapse and to inform any necessary post-remission treatment. A number of methodologies have been used to detect MRD in patients with hematologic malignancies, including flow cytometry, PCR-based assays, and next generation sequencing (NGS) tests.

Evaluation for MRD in bone marrow aspirate from patients with acute lymphoblastic leukemia (ALL) has demonstrated clinical utility; MRD testing has prognostic importance in predicting relapse and can help stratify high-risk patients in whom treatment intensification would be warranted from low-risk patients in whom such treatment (e.g., hematopoietic stem cell transplantation) could be avoided (Berry et al. 2017; Heikamp and Pui 2018; Kansagra et al. 2019; Eckert et al. 2019; Shah et al. 2020). Consensus recommendations indicate MRD assessments should be done in adults with ALL on first line treatment at various intervals and in relapsed or refractory ALL patients receiving salvage therapy. It is a vital component in the management of children and adults with ALL because of the association between risk for relapse and minimal residual disease (Berry et al. 2017).

The European LeukemiaNet (ELN) working party for MRD consisting of 24 experts from Europe and the United States published a consensus document in 2018 which provides recommendations to standardize and improve the reporting of MRD results. This group also provided clinical recommendations that MRD monitoring be considered part of the standard of care for all acute myeloid leukemia (AML) patients, but that molecular methods only be used for patients with subtypes

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

amenable to targeted **PCR-based** assays (specifically: APL, CBF AML, and NPM1-mutated AML). For others, flow cytometry is recommended (Schoorhuis et al. 2018).

While there is much hope that peripheral blood samples may be used for diagnosis and MRD detection in multiple myeloma (MM) in order to avoid the need for invasive biopsy, there are still many questions and technological hurdles to overcome (Soekojo et al. 2018; Romano et al. 2019; Mina et al. 2020). Intra-tumor heterogeneity adds to the complexities of detecting MRD with molecular testing. It is important to note that multiparametric flow cytometry (MFC) and NGS have not been directly compared, nor has NGS MRD testing been uniformly measured and reported in clinical trials (Bal et al. 2021). Presently, MRD results are not incorporated in treatment change decisions, and are mainly used as a prognostic measure (Rajkumar 2020; Bal et al. 2021).

Tumor Agnostic Testing

In recent years, there has been a great deal of progress in the development of targeted treatments for many types of cancer. Targeted therapies rely on the identification of the genetic mutations within tumor cells that drive the uncontrolled growth and proliferation of the cancer cells. These anticancer drugs interfere with and block the function of these specific molecular pathways. Many drugs have been incorporated into standard practice for the treatment of tumors with specific mutations. Some examples include EGFR-tyrosine kinase inhibitors, which are used to treat non-small cell lung cancers with EGFR mutations, and imatinib, which targets the BCR-ABL fusion gene that is characteristic of chronic myelogenous leukemia. However, clinical challenges also remain as tumors can develop resistance to these therapies. Combination treatments that target multiple pathways can be a more effective treatment strategy (Morris and Kopetz 2013).

Historically, US Food and Drug Administration (FDA) approvals of cancer treatments have been specific to the anatomical site of the primary tumor development, even when different types of cancers exhibit the same somatic variants. However, given growing evidence that unrelated tumor types can have the same molecular variants driving cancer development, current research into cancer therapies has started to focus on treatment based on molecular variants rather than the location of the tumor (Flaherty, Le, and Lemery 2017). As of late 2018, the FDA has approved a small number of therapies as tumor agnostic, meaning the treatment can be administered based on specific biomarkers rather than tumor location.

Microsatellite Instability and Immune Checkpoint Inhibitors

Microsatellites are highly polymorphic DNA sequences involving repeats of one to several base pairs. They occur in both coding and non-coding regions. These regions are prone to errors during DNA replication, which are typically repaired by DNA mismatch repair (MMR) enzymes. Evidence has supported the use of MSI testing to predict the effect of immune checkpoint inhibitors such as anti-PD-L1 antibodies (Le et al. 2015). The gold standard for MSI testing is by PCR or immunohistochemistry (IHC). Many tumor types have shown dramatic responses to this type of therapy, including undifferentiated malignancies (Devereaux et al. 2018). A recent retrospective analysis of cancer patients in the United States estimated that up to 44% of patients would be eligible for immune checkpoint inhibitors based on current FDA approval criteria, while only 13% would exhibit a favorable response (Haslam and Prasad 2019). Further research is ongoing to evaluate the optimal selection criteria for immune checkpoint inhibitors and additional treatment combinations for various types of cancers.

NTRK Fusion Testing

NTRK fusions are oncogenetic drivers that stimulate tumor growth in a wide variety of solid tumors. These fusions occur in developing tumor cells, and result in constitutive activation of the TRK tyrosine kinase domain, which includes the NTRK1, NTRK2 and NTRK3 genes. The ETV6-NTRK3 fusion

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

oncogene appears to be the dominant fusion event and has been seen in multiple cancer types including secretory breast carcinoma, mammary analogue secretory carcinoma (MASC), congenital fibrosarcoma, congenital mesoblastic nephroma, and acute myeloid leukemia. NTRK1 fusions have also been observed in lung adenocarcinoma, intrahepatic cholangiocarcinoma, spitzoid neoplasms, glioblastoma, and pontine glioma. NTRK2 fusions appear to be the least common to date. Tumor types with the highest known incidence of NTRK fusions include spitzoid neoplasms, secretory breast carcinoma, MASC, papillary thyroid cancer, congenital mesoblastic nephroma, and congenital fibrosarcoma. In most tumor types, NTRK fusions will only represent a small percentage of patients, if any. However, limitations in current testing methodologies make the true incidence of these fusions unknown (Vaishnavi et al. 2015).

The FDA has granted accelerated approval for larotrectinib (Vitrakvi) a small-molecule inhibitor of the tropomyosin receptor kinases that are encoded by NTRK genes. Due to the high degree of similarity between the NTRK genes larotrectinib is able to target all three (Yan and Zhang 2018). In August 2019, the FDA approved a second tumor agnostic medication, entrectinib (Rozlytrek) (AACR, 2019). These drugs are indicated for adult and pediatric patients with solid tumors positive for an NTRK gene fusion. These patients should have no known acquired resistance P/LP variant, and they must have metastatic disease or an unresectable tumor where the risk of surgery is high, and no other alternative therapeutic options exist.

Lung Cancer

A number of genetic changes within NSCLC tumors have been associated with improved response to various therapies, and best practice guidelines recommend molecular testing of advanced stage lung tumors, especially NSCLC adenocarcinomas, in order to help guide therapeutic decision-making. Epidermal growth factor receptor (EGFR) P/LP variant status has been shown to be significantly associated with tumor response to EGFR tyrosine kinase inhibitors (Lynch et al. 2004; Mok et al. 2009). This has led to the routine assessment of the presence of EGFR P/LP variants in advanced non-small cell lung cancers (NSCLC), particularly adenocarcinomas (Li et al. 2019). More recently, testing for EGFR pathogenic variants has also been shown to have clinical utility in the non-metastatic setting, specifically stages IB-IIIa (Wu et al. 2020). It is important to note that not all EGFR pathogenic variants have the same effect. For example, the p.T790M EGFR pathogenic variant is associated with relapse or resistance to TKI therapy. With the use of newer next generation sequencing assays, additional EGFR pathogenic variants are increasingly being identified in these patients, but there is limited data about the clinical implications of other types of EGFR pathogenic variants (Li et al. 2019).

Beyond EGFR, a number of additional genes may provide information about ideal treatment strategy or prognosis for patients with NSCLC. KRAS P/LP variants have been associated with primary EGFR TKI resistance as well as poor survival. Anaplastic lymphoma kinase (ALK) and ROS1 gene rearrangements have been identified in a subset of patients with NSCLC and are useful to identify patients for whom ALK or ROS1 inhibitors may be a very effective treatment strategy.

A number of other genetic alterations have been identified in individuals with NSCLC for which targeted therapies have already been developed for other tumor types, including: BRAF V600 P/LP variants, HER2 (ERBB2) P/LP variants, RET gene rearrangements, and MET amplification (Gregg et al. 2019). Multi-gene panel testing that includes these additional genes should be considered to identify patients who may benefit from targeted treatment (Lindeman et al. 2018).

Guidelines and recommendations regarding molecular testing in NSCLC tumor have been published by multiple societies including the American Society of Clinical Oncologists (ASCO), College of American Pathologists (CAP), the International Association for the Study of Lung Cancer (IASLC), and the Association for Molecular Pathology (AMP) (Lindeman et al. 2018; Hanna et al. 2017; Hanna et al. 2020; Kalemkerian et al. 2018). Based on high quality evidence, these groups agree with a strong recommendation that testing for ROS1, ALK, and EGFR mutations should be performed for all patients with advanced-stage (stages III B and above) lung adenocarcinoma. There is also agreement that testing for other genes, including BRAF, RET, ERBB2 (HER2), KRAS, MET, NTRK fusions, and PD-L1 amplification are also appropriate to aid in treatment decision-making in NSCLC, including tumors with histologies other than adenocarcinoma such as large cell or squamous cell carcinomas. In general, next generation sequencing panels are preferred, given the ability to analyze multiple genes from a single sample type, and to detect gene fusions/rearrangements and copy number alterations. Testing for P/LP variants within genes beyond those described above have not been incorporated into standard practice. Molecular testing for early-stage tumors, with the exception of EGFR for resected stage IB-IIIA tumors, is not included in these recommendations, given that these patients may be surgically cured with no need for molecularly targeted therapies (Lindeman et al. 2018; Hanna et al. 2017; Kalemkerian et al. 2018). Evaluation of tumor mutational burden has been proposed as an emerging biomarker to assess treatment response, however, there is no current consensus on how to measure this (Cyriac and Gandhi 2018).

While there has been success in broad molecular profiling and targeted therapies for NSCLC, there is a lack of evidence to support tumor testing for patients diagnosed with small cell lung cancer (SCLC) (Byers and Rudin 2015). Attempts to identify common driver P/LP variants in SCLC have revealed significant genetic heterogeneity across patients. The TP53 and RB1 genes are almost universally inactive in SCLC tumors, but targeted therapies for these genetic alterations are not yet available (Zaman and Bivona 2018). To date, there have been limited advances in the treatment of SCLC and there are specific challenges in performing genomic analysis on SCLC tumors compared to NSCLC tumors. Genomic profiling is currently being evaluated as an option, but more research is needed to demonstrate its effectiveness in this population (Umemura et al. 2015; Zaman and Bivona 2018).

Cancer Screening

Indeterminate Thyroid Nodules

Thyroid nodules occur in 1% of men and 5% of women (Haugen et al. 2016). These nodules are typically benign, although a small subset is malignant and require surgical resection with potential additional treatment. Cytological examination of FNA samples is the current standard of care for classifying thyroid nodules as malignant (thyroid carcinoma) or benign (thyroid adenoma), but this distinction is not always straightforward. Approximately 20-25% of samples are deemed indeterminate thyroid nodules (ITN) after being classified as Bethesda category III (atypia of undetermined significance/follicular lesion of undetermined significance, AUS/FLUS) or Bethesda category IV (follicular neoplasm/suspicious for a follicular neoplasm, FN/SFN). There are caveats that add complexity to ITN classification. The first is that approximately 10% of all FNA samples contain a significant Hurthle cell population. The second caveat came in early 2017, when the American Thyroid Association recommended a change in nomenclature from follicular variant of papillary thyroid carcinoma (FVPTC) to noninvasive follicular thyroid neoplasm with papillary-like nuclear features (NIFTP) in a subset of FNA with certain noninvasive features (Haugen et al. 2017). This move was based on evidence that these noninvasive tumors were indolent compared to infiltrative FVPTC and

could be managed in a much less aggressive manner by the avoidance of classifying this low-grade tumor as a carcinoma.

Traditionally, diagnostic surgery was performed for clarification and management of ITNs, but most procedures turned out to be unnecessary after data revealed up to 75% of cases were actually benign (Haugen et al. 2016). There is growing evidence that molecular diagnostic testing can alleviate the burden of surgical dependence in the reclassification of these indeterminate lesions for prognosis and treatment.

Gene expression classifiers (GECs) evaluate levels of RNA or miRNA expression to better understand gene regulation behavior. This can be important in predicting an abnormal pathological process, such as neoplastic growth. Genes included in these profiles may be proprietary and vary by laboratory. GECs used for ITN have a relatively low PPV and are generally considered “rule out” tests. An NPV of 95% is generally considered an acceptable threshold for this type of “rule out” test since the historical approach to observing nodules deemed cytologically benign left patients with a residual risk of 1-5% for malignancy (Ali et al. 2019). An abnormal result is not necessarily predictive of cancer, but if expression is normal, there is a high chance that cancer is currently not present. Long term data on the impact of conservative (observational) management for individuals with ITN and negative GEC results are still pending and are needed to fully establish clinical utility of GECs. In addition, there is currently insufficient independent prospective validation of performance characteristics of gene expression classifiers in Hurthle cells with available data encumbered by study limitations (e.g., low numbers and/or wide confidence intervals).

Tests that use next generation sequencing, point mutation analysis, or other targeted analyses of genes and P/LP variants known to have a strong association with thyroid malignancy (eg, BRAF, RET/PTC, RAS, PAX8/PPAR) are generally used as “rule in” tests. If a P/LP variant is identified, there is assumed to be a high likelihood that the thyroid nodule is malignant and requires surgical intervention. The prevalence of malignancy varies by the specific P/LP variant identified (Cohen et al. 2019), and the exact PPVs associated with these tests are highly variable.

Several professional societies have published guidelines regarding the use of molecular testing for indeterminate thyroid nodules and how to incorporate results into the management plan for patients with indeterminate cytology. The American Association of Clinical Endocrinologists do not recommend either in favor of or against the use of GECs for indeterminate thyroid nodules, due to insufficient evidence and limited follow-up. Molecular testing should not replace cytologic evaluation and should be considered when results are expected to influence clinical management. As a general rule molecular testing should not be considered in nodules with established benign or malignant cytologic characteristics (Gharib et al. 2016). Cytopathology expertise, patient characteristics and prevalence of malignancy within the population being tested impact NPV and PPV for molecular testing, but they do recommend it for BRAF and RET/PTC along with possibly PAX/PPARG and RAS P/LP variants if such detection is available (Gharib et al. 2016). With the exception of pathogenic variants such as BRAF V600E with PPV approaching 100% for PTC, evidence is insufficient to recommend in favor or against P/LP variant testing as a guide to determine the extent of surgery. Close follow-up is also still recommended for mutation-negative nodules or nodules classified as benign by a GEC because experience and follow-up for these is insufficient (Gharib et al. 2016).

The American Thyroid Association (ATA) issued a statement in 2015 regarding the surgical application of molecular profiling for thyroid nodules (Ferris et al. 2015). They suggest that a role exists for both molecular tumor profiling and gene expression classifier (GEC) systems in assisting with the

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

appropriate management of cytologically indeterminate nodules; however, the type of test chosen may be dependent upon additional clinical and sonographic features. They note that GECs may perform better when the initial suspicion for cancer is low, such as when the cytologic category is Bethesda III (AUS/FLUS), and that molecular testing performs better in settings with higher cancer frequencies (Haugen et al. 2016).

Prostate Cancer Early Screening

Prostate cancer is a common malignancy in men, and the worldwide burden of this disease is rising. Early detection and screening for prostate cancer is a clinical challenge, given the indolent nature of many prostate tumors as well as the risks and costs associated with overdiagnosis and overtreatment of this condition. Screening with prostate-specific antigen (PSA) was originally approved by the FDA in 1994, however, this method is controversial due to its low specificity and high rates of false positive results (Alford et al. 2017; Moyer 2012; Pinsky et al. 2017). Given the limitations of PSA screening, there is a clinical need for other methods to detect high-risk prostate cancers in the general population. Changes in the PSA threshold, frequency of screening, and the use of adjuvant tests (e.g., gene expression classifiers, digital rectal exam, mpMRI) have the potential to minimize the overdiagnosis and unnecessary biopsies associated with PSA screening. However, the best use of these options has yet to be established.

There are a number of genomic biomarker tests (e.g., PCA3, ConfirmMDx, ExoDx, SelectMDx) that have emerged in recent years with the goal of providing a more accurate method to aid in early detection of prostate cancer. PCA3 is a non-coding prostate-specific mRNA that is highly over-expressed in prostate cancer cells (median 66-fold up-regulation compared to adjacent benign tissue). The FDA has approved the use of this test for men age 50 or older who have had one or more previous negative prostate biopsies and for whom a repeat biopsy would be recommended by a urologist based on current standard of care (Narayan et al. 2017). The ConfirmMDx test is an epigenetic assay that evaluates the methylation status of the GTSP1, APC and RASSF1 genes. Methylation of these three genes can lead to a "field effect" which can indicate cancer nearby even if it was not included directly in the biopsy (Narayan et al. 2017). This test is most useful for deciding on repeat biopsy if PSA is high and initial biopsy is negative. ExoDx Prostate is a gene signature assay that evaluates expression of three genes known to play a role in prostate cancer initiation and progression: ERG, PCA3, and SPDEF. This test is considered a "rule-out" assay, as low-risk results indicate a low risk for high-grade prostate cancer and can support the decision to forego initial biopsy (McKiernan et al. 2016; McKiernan et al. 2018). SelectMDx measures expression levels of DLX1 and HOXC6 mRNA. Higher levels may be associated with an increased probability that prostate cancer will be detected on biopsy and increased risk of high-grade (Gleason score greater or equal to 7) prostate cancer, thus the test is intended to identify low risk patients who can safely avoid biopsy and proceed with active surveillance. Haese et al. (2019) concluded that the assay was optimized for biopsy native patients with serum PSA less than 10 ng/ml after clinical validation in 1955 men in a multicenter study.

The intended use of most gene expression classifier tests is to distinguish prostate cancer from benign prostatic conditions when a higher chance for cancer is suspected and many appear to have better sensitivity and specificity than PSA. However, results from gene expression profiles should not be interpreted as either positive or negative- instead, risk scores should be considered in the context of other tumor features (Cucchiara et al. 2018).

Professional Society Guidelines

American Cancer Society (ACS)

ACS 2018 Guideline Update. Colorectal cancer screening for average-risk adults.

Wolf AMD, Fontham ETH, Church TR, et al. CA Cancer J Clin. 2018 Jul;68(4):250-281. PubMed PMID: 29846947.

American College of Medical Genetics and Genomics (ACMG)

ACMG Statement. Points to Consider for Reporting of Germline Variation in Patients Undergoing Tumor Testing.

Li M.M., Chao E., Espalin E.D. et al. Genet Med. 2020 Apr 23. PubMed PMID: 32321997.

American Society of Clinical Oncology (ASCO)

ASCO Clinical Practice Guideline Endorsement Update Summary. Role of patient and disease factors in adjuvant systemic therapy decision making for early-stage, operable breast cancer

Henry NL, Somerfield MR, Abramson VG, et al. Update of the ASCO Endorsement of the Cancer Care Ontario Guideline. J Clin Oncol. 2019 Aug 1;37(22):1965-1977. PubMed PMID: 31206315.

ASCO Clinical Practice Guideline. Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women with Early-Stage Invasive Breast Cancer.

2019 Update Summary. Andre F, Ismaila N, Stearns V. J Oncol Pract. 2019 Sep;15(9):495-497. PubMed PMID: 31306037. Original Publication. Harris LN, Ismaila N, McShane LM, et al. J Clin Oncol. 2016 Apr 1;34(10):1134-50. PubMed PMID: 26858339.

ASCO Clinical Practice Guideline. Neoadjuvant Chemotherapy, Endocrine Therapy, and Targeted Therapy for Breast Cancer.

Krode LA, Somerfield MR, Carey LA. J Clin Oncol. 2021 Jan 28;JCO2003399. PubMed PMID: 33507815.

ASCO Clinical Practice Guideline. Management of Male Breast Cancer

Hassett MJ, Somerfield MR, Baker ER, et al. J Clin Oncol. 2020 Feb 14;JCO1903120. PubMed PMID: 32058842.

ASCO Clinical Practice Guideline. Molecular Biomarkers in Localized Prostate Cancer.

Egger SE, Rumble RB, Armstrong AJ, et al. J Clin Oncol. 2020;38(13):1474-1494. PubMed PMID: 31829902.

ASCO Guideline Endorsement. Molecular Testing for the Selection of Patients with Lung Cancer for Treatment with Targeted Tyrosine Kinase Inhibitors.

Kalemkerian GP, Narula N, Kennedy EB, et al. ASCO Endorsement of the College of American Pathologists/International Association for the Study of Lung Cancer/Association for Molecular Pathology Clinical Practice Guideline Update. J Clin Oncol. 2018 Mar 20;36(9):911-919. PubMed PMID: 29401004.

ASCO Guideline. Therapy for Stage IV Non-Small-Cell Lung Cancer Without Driver Alterations.

Hanna N, Schneider B, Temin S, et al. Joint Guideline Update by ASCO and the Ontario Health (OH) Cancer Care Ontario (CCO). J Clin Oncol. 2020 May 10;38(14):1608-1632. PubMed PMID: 31990617.

American Thyroid Association (ATA)

ATA Surgical Statement. Surgical Application of Molecular Profiling for Thyroid Nodules: Current Impact on Perioperative Decision Making.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Ferris RL, Baloch Z, Bernet V, et al. *Thyroid*. 2015 Jul;25(7):760-8. PubMed PMID: 26058403.

ATA Guideline. Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer.

Haugen BR, Alexander EK, Bible KC, et al. *ATA Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer*. 2015 *Thyroid*. 2016 Jan;26(1):1-133. Epub 2017 Feb 21. PubMed PMID: 28114862.

ATA Guideline. Management of Thyroid Nodules and Differentiated Thyroid Cancer. Review and Recommendation on the Proposed Renaming of Encapsulated Follicular Variant Papillary Thyroid Carcinoma Without Invasion to Noninvasive Follicular Thyroid Neoplasm with Papillary-Like Nuclear Features.

Haugen BR, Sawka AM, Alexander EK, et al. *Thyroid*. 2017 Apr;27(4):481-3. Epub 2017 Feb 21. PubMed PMID: 28114862.

European Association of Urology (EAU)

Biochemical Recurrence in Prostate Cancer: The European Association of Urology Prostate Cancer Guidelines Panel Recommendations.

Van den Broeck T, et al. *Eur Urol Focus*. 2020 Mar 15;6(2):231-234. PubMed PMID: 31248850.

European Society for Medical Oncology

Cancers of Unknown Primary Site: ESMO Clinical Practice Guidelines for Diagnosis, Treatment and Follow-up.

Fizazi K, Greco FA, Pavlidis N, et al. *Ann Oncol*. 2015 Sep;26 Suppl 5:v133-8. PubMed PMID: 26314775.

ESMO Recommendations on Predictive Biomarker Testing for Homologous Recombination Deficiency and PARP Inhibitor Benefit in Ovarian Cancer.

Miller RE, Leary A, Scott C, et al. *Ann Oncol*. 2020 Dec;31(12):1606-1622. PubMed PMID: 33004253.

Prostate Cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up.

Parker C, Castro E, Fizazi K, et al.; ESMO Guidelines Committee. *Ann Oncol*. 2020 Jun 25;S0923-7534(20)39898-7. PubMed PMID: 32593798.

Society of Gynecologic Oncology

Endometrial Cancer: A Society of Gynecologic Oncology Evidence-Based Review and Recommendations

Hamilton C, Pothuri B, Arend R, et al. *Gynecol Oncol*. 2021 Jan 27;S0090-8258(20)34224-4. PubMed PMID: 33516529.

St. Gallen Consensus Conference (Vienna)

Optimal Primary Breast Cancer Treatment. Brief Summary of Consensus Discussion.

Balic M, Thomssen C, Würtle R, et al: *St. Gallen/Vienna 2019. Breast Care* 2019;14:103-110. PubMed PMID: 31798382.

US Preventive Services Task Force (USPSTF)

USPSTF Recommendation Statement. Screening for Colorectal Cancer.

JAMA. 2016;315(23):2564-75. PubMed PMID: 27304597.

World Health Organization (WHO)

WHO Classification of Myeloid Neoplasms and Acute Leukemia (2016 Revision).

Arber DA, Orazi A, Hasserjian R et al. *Blood*. 2016 May 19;127(20):2391-405. PubMed PMID: 27069254.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Joint Statements

Circulating Tumor DNA Analysis in Patients with Cancer: ASCO and CAP Joint Review.

Merker JD, Oxnard GR, Compton C, et al. *J Clin Oncol*. 2018 Jun 1;36(16):1631-1641. PubMed PMID: 29504847.

Clinically localized prostate cancer: AUA/ASTRO/SUA Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options.

Sanda, Martin G. et al. *J Urol*. 2018. Mar; 199(3): 683-690. PubMed PMID: 29203269.

Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part II: Recommended approaches and details of specific care options.

Sanda MG, Cadeddu JA, Kirkby E, et al. *J Urol*. 2018 Jan 10. PubMed PMID: 29331546.

Colorectal Cancer Screening: Recommendations for Physicians and Patients from the U.S. Multi-Society Task Force on Colorectal Cancer.

Rex DK, Boland CR, Dominitz J. *Am J Gastroenterol*. 2017 Jul;112(7):1016-30. PubMed PMID: 28555630.

Minimal/Measurable Residual Disease in AML: A Consensus Document from the European LeukemiaNet MRD Working Party.

Schuurhuis GJ, Heuser M, Freeman S, et al. *Blood*. 2018 Mar 22;131(12):1275-1291. PubMed PMID: 29330221.

Molecular Biomarkers for the Evaluation of Colorectal Cancer: Guideline from the American Society for Clinical Pathology, College of American Pathologists, Association for Molecular Pathology, and American Society of Clinical Oncology.

Sepulveda AR, Hamilton SR, Allegra CJ, Grody W, Cushman-Vokoun AM, et al. *J Mol Diagn*. 2017 Mar;19(2):187-225. PubMed PMID: 28185757.

Prostate Cancer. Guideline from the EAU, EANM, ESTRO, ESUR, SIOG.

Mottet N, Cornford P, van den Bergh E., et al. 2020. Available at <https://uroweb.org/guideline/prostate-cancer/>. Accessed on 1/28/2021.

Updated Molecular Testing Guideline for the Selection of Lung Cancer Patients for Treatment with Targeted Tyrosine Kinase Inhibitors. Guideline from the College of American Pathologists, the International Association for the Study of Lung Cancer, and the Association for Molecular Pathology.

Lindeman NI, Cagle PT, Aisner DL, et al. *J Mol Diagn*. 2018 Mar;20(2):129-159. PubMed PMID: 29398453.

Selected References

- 1 Abida W, Cyrta J, Heller G, et al. Genomic correlates of clinical outcome in advanced prostate cancer. *Proc Natl Acad Sci U S A*. 2019 May 6. pii: 201902651. Doi: 10.1073/pnas.1902651116. [Epub ahead of print] PubMed PMID: 31061129.
- 2 Albert CM, Davis JL, Federman N, et al. TRK fusion cancers in children: A clinical review and recommendations for screening. *J Clin Oncol*. 2019 Feb 20;37(6):513-524. PubMed PMID: 30592640.
- 3 Alghasham N, Alnouri Y, Abalkhil H, et al. Detection of mutations in JAK2 exons 12-15 by Sanger sequencing. *Int J Lab Hematol*. 2016 Feb;38(1):34-41. Epub 2015 Sep 11. PubMed PMID: 26361084.
- 4 Albain KS, Barlow WE, Shak S, et al. Prognostic and predictive value of the 21-gene recurrence score assay in postmenopausal women with node-positive, estrogen-receptor-positive breast cancer on chemotherapy: a retrospective analysis of a randomised trial. *Lancet Oncol*. 2010 Jan;11(1):55-65. Epub 2009 Dec 10. PubMed PMID: 20005174.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 5 Alexander EK, Schorr M, Klopffer J, et al. Multicenter clinical experience with the Afirma Gene Expression Classifier. *J Clin Endocrinol Metab.* 2014 Jan;99(1):119-25. Epub 2013 Dec 20. PubMed PMID: 24152684.
- 6 Alford AV, Brito JM, Yadav KK, Yadav SS, Tewari AK, Renzulli J. The Use of Biomarkers in Prostate Cancer Screening and Treatment. *Reviews in Urology.* 2017;19(4):221-234. PubMed PMID: 29472826.
- 7 Andre F, Ciruelos E, Rubovsky G, Campone M, Loibl S, Rugo HS, Iwata H, Conte P, Mayer IA, Kaufman B, Yamashita T, Lu YS, Inoue K, Takahashi M, Papai Z, Longin AS, Mills D, Wilke C, Hirawat S, Juric D, SOLAR-1 Study Group. Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer. *N Engl J Med.* 2019 May 16;380(20):1929-1940. PubMed PMID: 31091374.
- 8 Armstrong AJ, Halabi S, Luo J, et al. Prospective multicenter validation of androgen receptor splice variant 7 and hormone therapy resistance in high-risk castration-resistant prostate cancer: The PROPHECY Study. *J Clin Oncol.* 2019 May 1;37(13):1120-1129. doi: 10.1200/JCO.18.01731. Epub 2019 Mar 13. PubMed PMID: 30865549.
- 9 Arpino G, Generali D, Sapino A, et al. Gene expression profiling in breast cancer: A clinical perspective. *Breast.* 2013 Apr;22(2):109-20. Epub 2013 Feb 23. PubMed PMID: 23462680.
- 10 Ayala de la Peña F, Andrés R, García-Sáenz JA, et al. SEOM clinical guidelines in early stage breast cancer (2018). *Clin Transl Oncol.* 2019 Jan;21(1):18-30. PMID: 30443868.
- 11 Azim HA Jr, Michiels S, Zagouri F, et al. Utility of prognostic genomic tests in breast cancer practice: The IMPAKT 2012 Working Group Consensus Statement. *Ann Oncol.* 2013;24(3):647-54. Epub 2013 Jan 20. PubMed PMID: 23337633.
- 12 Bal S, Giri S, Godby KN, Costa LJ. New regimens and directions in the management of newly diagnosed multiple myeloma. *Am J Hematol.* 2021. PubMed PMID: 33393136.
- 13 Berry DA, Zhou S, Higley H, Mukundan L, Fu S, Reaman GH, Wood BL, Kelloff GJ, Jessup JM, Radich JP. Association of minimal residual disease with clinical outcome in pediatric and adult acute lymphoblastic leukemia: a meta-analysis. *JAMA Oncol.* 2017 Jul 13;3(7). PubMed PMID: 28494052.
- 14 Binder C, Matthes KL, Korol D, Rohrmann S, Moch H. Cancer of unknown primary-Epidemiological trends and relevance of comprehensive genomic profiling. *Cancer Med.* 2018 Sep;7(9):4814-4824. PubMed PMID: 30019510.
- 15 Blok EJ, Bastiaannet E, van den Hout WB, Liefers GJ, Smit VTHBM, Kroep JR, van de Velde CJH. Systematic review of the clinical and economic value of gene expression profiles for invasive early breast cancer available in Europe. *Cancer Treat Rev.* 2018 Jan; 62:74-90. Review. PMID: 29175678.
- 16 Bombard Y, Bach PB, Offit K. Translating genomics in cancer care. *J Natl Compr Canc Netw.* 2013 Nov;11(11):1343-53. PubMed PMID: 24225968.
- 17 Buus R, Sestak I, Kronenwett R, et al. Comparison of EndoPredict and EPclin with Oncotype DX recurrence score for prediction of risk of distant recurrence after endocrine therapy. *J Natl Cancer Inst.* 2016 Jul 10;108(11). PubMed PMID: 27400969.
- 18 Byers LA, Rudin CM. Small cell lung cancer: where do we go from here? *Cancer.* 2015 Mar 1;121(5):664-72. PubMed PMID: 25336938.
- 19 Cao SS, Lu CT. Recent perspectives of breast cancer prognosis and predictive factors. *Oncol Lett.* 2016 Nov;12(5):3674-3678. PMID: 27900052.
- 20 Cardoso F, van't Veer LJ, Bogaerts J, et al. 70-Gene Signature as an aid to treatment decisions in early-stage breast cancer. *N Engl J Med.* 2016 Aug 25; 375(8):717-29. PubMed PMID: 27557300.
- 21 Chang MC, Souter LH, Kamel-Reid S, et al; Molecular Oncology Advisory Committee. Clinical utility of multigene profiling assays in early-stage breast cancer. *Current oncology (Toronto, Ont.)* vol. 24,5 (2017): e403-e422. *Curr Oncol.* 2017 Oct;24(5): e403-e422. PubMed PMID: 29089811.
- 22 Chia SKL. Clinical application and utility of genomic assays in early-stage breast cancer: key lessons learned to date. *Curr Oncol.* 2018 Jun;25(Suppl 1):S125-S130. Doi: 10.3747/co.25.3814. Epub 2018 Jun 13. Review. PMID: 29910655.
- 23 Colomer R, Aranda-López I, Albanell J, et al. Biomarkers in breast cancer: A consensus statement by the Spanish Society of Medical Oncology and the Spanish Society of Pathology. *Clin Transl Oncol.* 2018 Jul;20(7):815-826. PubMed PMID: 29273958.
- 24 Cucchiara V, Cooperberg MR, Dall'Era M, Lin DW, Montorsi F, Schalken JA, Evans CP. Genomic Markers in Prostate Cancer Decision Making. *Eur Urol.* 2018 Apr;73(4):572-582. PubMed PMID: 29129398.
- 25 Curigliano G, Burstein HJ, P Winer E, et al. De-escalating and escalating treatments for early-stage breast cancer: the St. Gallen International Expert Consensus Conference on the Primary Therapy of Early Breast Cancer 2017. *Ann Oncol.* 2017 Aug 1;28(8):1700-1712. PMID: 28838210.
- 26 Cuzick J, Thorat MA, Andriole G, et al. Prevention and early detection of prostate cancer. *Lancet Oncol.* 2014 Oct;15(11): e484-92. PubMed PMID: 25281467.
- 27 Cyriac G, Gandhi L. Emerging biomarkers for immune checkpoint inhibition in lung cancer. *Semin Cancer Biol.* 2018 Oct;52(Pt 2):269-277. PubMed PMID: 29782924.
- 28 Denkert C, Kronenwett R, Schlake W, et al. Decentral gene expression analysis for ER+/Her2- breast cancer: results of a proficiency testing program for the EndoPredict assay. *Virchows Arch.* 2012 Mar; 460(3):251-259. PubMed PMID: 22371223.
- 29 Devereaux KA, Charu V, Zhao S, Charville GW, Bangs CD, van de Rijn M, Cherry AM, Natkunam Y. Immune checkpoint blockade as a potential therapeutic strategy for undifferentiated malignancies. *Hum Pathol.* 2018 Dec;82:39-45. PubMed PMID: 30539796.
- 30 Douillard JY, Ostoros G, Cobo M, et al. Gefitinib treatment in EGFR mutated caucasian NSCLC: circulating, free tumor DNA as a surrogate for determination of EGFR status. *J Thorac Oncol* 2014; 9:1345-53. PubMed PMID: 25122430.
- 31 Dubsy P, Brase JC, Jakesz R, et al. The EndoPredict score provides prognostic information on late distant metastases in ER+/HER2- breast cancer patients. *Br J Cancer.* 2013;109(12):2959-2964. PubMed PMID: 24157828.
- 32 Dubsy P, Filipits M, Jakesz R, et al. EndoPredict improves the prognostic classification derived from common clinical guidelines in ER-positive, HER2-negative early breast cancer. *Ann Oncol.* 2013 Mar;24(3):640-647. PubMed PMID: 23035151.
- 33 Duffy MJ, Harbeck N, Nap M, et al. Clinical use of biomarkers in breast cancer: updated guidelines from the European Group on Tumor Markers (EGTM). *Eur J Cancer.* 2017 Apr; 75:284-298. Epub 2017 Feb 28. PubMed PMID: 28259011.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 34 Duick D, Klopper J, Diggans J, et al. The impact of benign gene expression classifier test results on the endocrinologist-patient decision to operate on patients with thyroid nodules with indeterminate fine-needle aspiration cytopathology. *Thyroid*. Oct 2012; 22(10): 996–1001. Epub 2012 Aug 8. PubMed PMID: 22873825.
- 35 Engelman JA, Chen L, Tan X, et al. Effective use of PI3K and MEK inhibitors to treat mutant Kras G12D and PIK3CA H1047R murine lung cancers. *Nat Med*. 2008 Dec;14(12):1351-6. Epub 2008 Nov 30. PubMed PMID: 19029981.
- 36 Fiala O, Pesek M, Finek J, et al. Gene mutations in squamous cell NSCLC: insignificance of EGFR, KRAS and PIK3CA mutations in prediction of EGFR-TKI treatment efficacy. *Anticancer Res*. 2013 Apr;33(4):1705-11. PubMed PMID: 23564819.
- 37 Fidler MJ, Morrison LE, Basu S, et al. PTEN and PIK3CA gene copy numbers and poor outcomes in non-small cell lung cancer patients with gefitinib therapy. *Br J Cancer*. 2011 Dec 6;105(12):1920-6. Epub 2011 Nov 17. PubMed PMID: 22095222.
- 38 Filipits M, Rudas M, Jakesz R, et al. A new molecular predictor of distant recurrence in ER-positive, HER2-negative breast cancer adds independent information to conventional clinical risk factors. *Clin Cancer Res* 2011; 17:6012-6020. PubMed PMID: 21807638.
- 39 Flaherty KT, Le DT, Lemery S. Tissue-Agnostic Drug Development. *Am Soc Clin Oncol Educ Book*. 2017; 37:222-230. PubMed PMID: 28561648.
- 40 Gokbuget N, Dombret H, Bonifacio M, et al. Binatumomab for minimal residual disease in adults with B-cell precursor acute lymphoblastic leukemia. *Blood*. 2018; 131(14):1522-1531. PubMed PMID: 29358182.
- 41 Goncalves MD, Hopkins BD, Cantley LC. Phosphatidylinositol 3-kinase, growth disorders, and cancer. *N Engl J Med*. 2018 Nov 22;379(21):2052-2062. doi: 10.1056/NEJMra1704560. Review. PubMed PMID: 30462943.
- 42 Gore JL, du Plessis M, Santiago-Jimenez M, et al. Decipher test impacts decision making among patients considering adjuvant and salvage treatment after radical prostatectomy: Interim results from the Multicenter Prospective PRO-IMPACT study. *Cancer*. 2017 Aug 1;123(15):2850-9. Epub 2017 Apr 19. PubMed PMID: 28422278.
- 43 Grinfeld J, Nangalia J, Baxter EJ, et al. Classification and personalized prognosis in myeloproliferative neoplasms. *N Engl J Med*. 2018 Oct 11; 379(15):1416-1430. PubMed PMID: 30304655.
- 44 Haese A, Trooskens G, Steyaert S, et al. Multicenter Optimization and Validation of a 2-Gene mRNA Urine Test for Detection of Clinically Significant Prostate Cancer Prior to Initial Prostate Biopsy. *J Urol*. 2019 PubMed PMID: 31026217.
- 45 Hang JF, Westra WH, Cooper DS, Ali SZ. The impact of noninvasive follicular thyroid neoplasm with papillary-like nuclear features on the performance of the Afirma gene expression classifier. *Cancer*. 2017 Sep; 125(9):683-91. Epub 2017 May 24. PubMed PMID: 28544601.
- 46 Hanna N, Johnson D, Temin S, et al. Systemic Therapy for Stage IV Non-Small-Cell Lung Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol*. 2017 Oct 20;35(30):3484-3515. Epub 2017 Aug 14. PubMed PMID: 28806116.
- 47 Harnan S, Tappenden P, Cooper K, et al. Tumor profiling tests to guide adjuvant chemotherapy decisions in early breast cancer: a systemic review and economic analysis. *Health Technol Assess*. 2019 Jun;23(30):1-328. PubMed PMID: 31264581.
- 48 Harris LN, Ismaila N, McShane LM, et al. Use of biomarkers to guide decisions on adjuvant systemic therapy for women with early-stage invasive breast cancer: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2016 Feb 8. Pii: JCO652289. [Epub ahead of print] PubMed PMID: 26858339.
- 49 Hayashi H, Kurata T, Takiguchi Y, Arai M, Takeda K, Akiyoshi K, Matsumoto K, Onoe T, Mukai H, Matsubara N, Minami H, Toyoda M, Onozawa Y, Ono A, Fujita Y, Sakai K, Koh Y, Takeuchi A, Ohashi Y, Nishio K, Nakagawa K. Randomized Phase II Trial Comparing Site-Specific Treatment Based on Gene Expression Profiling With Carboplatin and Paclitaxel for Patients With Cancer of Unknown Primary Site. *J Clin Oncol*. 2019 Mar 1;37(7):570-579. PubMed PMID: 30653423.
- 50 Hechtman JF, Ross DS. The past, present, and future of HER2 (ERBB2) in cancer: Approaches to molecular testing and an evolving role in targeted therapy. *Cancer Cytopathol*. 2019 Jul;127(7):428-431. PubMed PMID: 30938930.
- 51 Herlemann A, Washington SL 3rd, Eapen RS, Cooperberg MR. Whom to treat: postdiagnostic risk assessment with gleason score, risk models, and genomic classifier. *Urol Clin North Am*. 2017 Nov;44(4):547-555. PubMed PMID: 29107271.
- 52 Hong DS, Bauer TM, Lee JJ, et al. Larotrectinib in adult patients with solid tumours: a multi-centre, open-label, phase I dose-escalation study. *Ann Oncol*. 2019 Feb 1;30(2):325-331. PubMed PMID: 30624546.
- 53 Janku F, Huang HJ, Claes B, et al. BRAF mutation testing in cell-free DNA from the plasma of patients with advanced cancers using a rapid, automated molecular diagnostics system. *Mol Cancer Ther*. 2016 Jun;15(6):1397-404. Epub 2016 May 20. PubMed PMID: 27207774.
- 54 Kalinsky K, Barlow W, Meric-Bernstam F, et al. GS3-00. First results from a phase III randomized clinical trial of standard adjuvant endocrine therapy (ET) +/- chemotherapy (CT) in patients (pts) with 1-3 positive nodes, hormone receptor-positive (HR+) and HER2-negative (HER2-) breast cancer (BC) with recurrence score (RS) ≤25: SWOG S1007 (RxPonder). 2020 Dec 10: San Antonio Breast Cancer Symposium. <https://www.sabcs.org/2020-SABCS>.
- 55 Keedy VL, Temin S, Somerfield MR, et al. American Society of Clinical Oncology provisional clinical opinion: epidermal growth factor receptor (EGFR) mutation testing for patients with advanced non-small-cell lung cancer considering first-line EGFR tyrosine kinase inhibitor therapy. *J Clin Oncol*. 2011 May 20;29(15):2121-7. Epub 2011 Apr 11. PubMed PMID: 21482992.
- 56 Kloos RT, Reynolds JD, Walsh PS, et al. Does addition of BRAF V600E mutation testing modify sensitivity or specificity of the Afirma Gene Expression Classifier in cytologically indeterminate thyroid nodules? *J Clin Endocrinol Metab*. 2013 Apr;98(4): E761-8. Epub 2013 Mar 8. PubMed PMID: 23476074.
- 57 Kretschmer A, Tilki D. Biomarkers in prostate cancer - Current clinical utility and future perspectives. *Crit Rev Oncol Hematol*. 2017 Dec; 120:180-193. PubMed PMID: 29198331.
- 58 Kronenwett R, Bohmann K, Prinzler J, et al. Decentral gene expression analysis: analytical validation of the Endopredict genomic multianalyte breast cancer prognosis test. *BMC Cancer*. 2012 Oct 5; 12:456. PubMed PMID: 23039280.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 59 Krop I, Ismaila N, Andre F, et al. Use of biomarkers to guide decisions on adjuvant systemic therapy for women with early-stage invasive breast cancer: American Society of Clinical Oncology clinical practice guideline focused update. *J Clin Oncol* 2017; 35:2838-2847. PubMed PMID: 28692382.
- 60 Lamy PJ, Allory Y, Gauchez AS, et al. Prognostic Biomarkers Used for Localised Prostate Cancer Management: A Systematic Review. *Eur Urol Focus*. 2017 Mar 7. pii: S2405-4569(17)30065-2. PubMed PMID: 28753865.
- 61 Lasho TL, Mudireddy M, Finke CM, et al. Targeted next-generation sequencing in blast phase myeloproliferative neoplasms. *Blood Adv*. 2018 Feb 27;2(4):370-380. PubMed PMID: 29467191.
- 62 Le DT, Uram JN, Wang H, et al. PD-1 Blockade in Tumors with Mismatch-Repair Deficiency. *N Engl J Med*. 2015 Jun 25;372(26):2509-20. PubMed PMID: 26028255.
- 63 Levy B, Hu ZI, Cordova KN, Close S, et al. Clinical Utility of Liquid Diagnostic Platforms in Non-Small Cell Lung Cancer. *Oncologist*. 2016 Sep;21(9):1121-30. Review. PMID: 27388233.
- 64 Li Y, Appius A, Pattipaka T, Feyereislova A, Cassidy A, Ganti AK. Real-world management of patients with epidermal growth factor receptor (EGFR) mutation-positive non-small-cell lung cancer in the USA. *PLoS One*. 2019 Jan 4;14(1). PubMed PMID: 30608948.
- 65 Liebs S, Keilholz U, Kehler I, Schweiger C, Haybäck J, Nonnenmacher A. Detection of mutations in circulating cell-free DNA in relation to disease stage in colorectal cancer. *Cancer Med*. 2019 Jul;8(8):3761-3769. PubMed PMID: 31134762.
- 66 Lin DW, Nelson PS. Prognostic Genomic Biomarkers in Patients With Localized Prostate Cancer: Is Rising Utilization Justified by Evidence? *JAMA Oncol*. 2021 Jan 1;7(1):59-60. PubMed PMID: 33237305.
- 67 Livhits MJ, Zhu CY, Kuo EJ, Nguyen DT, Kim J, Tseng CH, Leung AM, Rao J, Levin M, Douek ML, Beckett KR, Cheung DS, Gofnung YA, Smooke-Praw S, Yeh MW. Effectiveness of Molecular Testing Techniques for Diagnosis of Indeterminate Thyroid Nodules: A Randomized Clinical Trial. *JAMA Oncol*. 2021 Jan 1;7(1):70-77. PubMed PMID: 33300952.
- 68 Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med*. 2004 May 20;350(21):2129-39. Epub 2004 Apr 29. PMID: 15118073.
- 69 Madison R, Schrock AB, Castellanos E, et al. Retrospective analysis of real-world data to determine clinical outcomes of patients with advanced non-small cell lung cancer following cell-free circulating tumor DNA genomic profiling. *Lung Cancer*. 2020 Aug 6;148:69-78. PubMed PMID: 32823229.
- 70 Martin M, Brase JC, Calvo L, et al. Clinical validation of the EndoPredict test in node-positive, chemotherapy-treated ER+/HER2- breast cancer patients: results from the GEICAM 9906 trial. *Breast Cancer Res*. 2014 Apr 12;16(2): R38. PubMed PMID: 24725534.
- 71 McClure RF, Ewalt MD, Crow J, et al. Clinical Significance of DNA Variants in Chronic Myeloid Neoplasms (CMNs): A Report of the Association for Molecular Pathology. *J Mol Diagn*. 2018 Aug 20. pii: S1525-1578(17)30409-9. PubMed PMID: 30138727.
- 72 McIver B, Castro MR, Morris JC, et al. An independent study of a gene expression classifier (Afirma) in the evaluation of cytologically indeterminate thyroid nodules. *J Clin Endocrinol Metab*. 2014 Nov;99(11):4069-77. Epub 2014 Apr 29. PubMed PMID: 24780044.
- 73 McKiernan J, Donovan MJ, O'Neill V, et al. A novel urine exosome gene expression assay to predict high-grade prostate cancer at initial biopsy. *JAMA Oncol*. 2016 Jul 1;2(7):882-889. PubMed PMID: 30237023.
- 74 McKiernan J, Donovan MJ, Margolis E, et al. A prospective adaptive utility trial to validate performance of a novel urine exosome gene expression assay to predict high-grade prostate cancer in patients with prostate-specific antigen 2-10ng/ml at Initial Biopsy. *Eur Urol*. 2018 Sep 17. PMID: 30237023.
- 75 Meleth S, Whitehead N, Evans TS, Lux L. Agency for Healthcare Research and Quality (US)Centers for Medicare & Medicaid (CMS) Agency for Healthcare Research and Quality (AHRQ) Technology Assessment on Genetic Testing or Molecular Pathology Testing of Cancers with Unknown Primary Site to Determine Origin; 2013 Feb. PubMed PMID: 25855837.
- 76 Mina R, Oliva S, Boccadoro M. Minimal Residual Disease in Multiple Myeloma: State of the Art and Future Perspectives. *J Clin Med*. 2020 Jul 7;9(7):E2142. PubMed PMID: 32645952.
- 77 Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med*. 2009 Sep 3;361(10):947-57. Epub 2009 Aug 19. PubMed PMID: 19692680.
- 78 Morris V, Kopetz S. BRAF inhibitors in clinical oncology. *F1000Prime Rep*. 2013 Apr 2;5:11. PubMed PMID: 23585929.
- 79 Moyer VA. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Annals of internal medicine* 2012; 157:120-134. PubMed PMID: 22801674.
- 80 Müller BM, Keil E, Lehmann A, et al. The EndoPredict Gene-Expression Assay in Clinical Practice - Performance and Impact on Clinical Decisions. *PLoS One*. 2013 Jun 27;8(6): e68252. PubMed PMID: 23826382.
- 81 Myint NNM, Verma AM, Fernandez-Garcia D, et al. Circulating tumor DNA in patients with colorectal adenomas: assessment of detectability and genetic heterogeneity. *Cell Death Dis*. 2018 Aug 30;9(9):894. PubMed PMID: 30166531.
- 82 Narayan VM, Konety BR, Warlick C. Novel biomarkers for prostate cancer: An evidence-based review for use in clinical practice. *Int J Urol*. 2017 May;24(5):352-360. Epub 2017 Mar 27. PubMed PMID: 28345187.
- 83 Nitz U, Gluz O, Christgen M, et al. Reducing chemotherapy use in clinically high-risk, genomically low-risk pN0 and pN1 early breast cancer patients: five-year data from the prospective, randomised phase 3 West German Study Group (WSG) PlanB trial. *Breast Cancer Res Treat*. 2017 Oct;165(3):573-583. Erratum in: *Breast Cancer Res Treat*. 2019 Jan 10. PubMed PMID: 28664507.
- 84 Paik S, Tang G, Shak S, et al. Gene expression and benefit of chemotherapy in women with node-negative, estrogen receptor-positive breast cancer. *J Clin Oncol*. 2006; 24:3726-3734. PubMed PMID: 16720680.
- 85 Parikh RB, Prasad V. Blood-based screening for colon cancer: A disruptive innovation or simply a disruption? *JAMA*. 2016 Jun 21;315(23):2519-20. PubMed PMID: 27305625.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 86 Pinsky PF, Prorok PC, Kramer BS. Prostate cancer screening - a perspective on the current state of the evidence. *N Engl J Med*. 2017 Mar 30;376(13):1285-9. PubMed PMID: 28355509.
- 87 Ragon BK, Savona MR. The challenge of treating myelodysplastic syndromes/myeloproliferative neoplasms. *Clin Lymphoma Myeloma Leuk*. 2017 Jul; 17S:S37-42. PubMed PMID: 28760301.
- 88 Rajkumar SV, Dimopoulos MA, Palumbo A, Blade J, Merlini G, Mateos MV, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol*. 2014 Nov; 15(12): e538-548. PubMed PMID: 25439696.
- 89 Rajkumar SV. Multiple myeloma: 2020 update on diagnosis, risk-stratification and management. *Am J Hematol*. 2020 May;95(5):548-567. Erratum in: *Am J Hematol*. 2020 Nov;95(11):1444. PubMed PMID: 32212178.
- 90 Ravandi F. How I treat Philadelphia chromosome-positive acute lymphoblastic leukemia. *Blood*. 2019 Jan 10;133(2):130-136. PubMed PMID: 30442680.
- 91 Rolfo C, Mack PC, Scagliotti GV, et al. Liquid Biopsy for Advanced Non-Small Cell Lung Cancer (NSCLC): A Statement Paper from the IASLC. *J Thorac Oncol*. 2018 Sep;13(9):1248-1268. PubMed PMID: 29885479.
- 92 Rolfo C, Cardona AF, Cristofanilli M, et al. Challenges and opportunities of cfDNA analysis implementation in clinical practice: Perspective of the International Society of Liquid Biopsy (ISLB). *Crit Rev Oncol Hematol*. 2020 Jul;151:102978. PubMed PMID: 32428812.
- 93 Romano A, et al. Minimal Residual Disease Assessment Within the Bone Marrow of Multiple Myeloma: A Review of Caveats, Clinical Significance and Future Perspectives. *Front Oncol*. 2019 Aug 20;9:699. PubMed PMID: 31482061.
- 94 Salto-Tellez M, Tsao MS, Shih JY, et al. Clinical and testing protocols for the analysis of epidermal growth factor receptor mutations in East Asian patients with non-small cell lung cancer: a combined clinical-molecular pathological approach. *J Thorac Oncol*. 2011 Oct;6(10):1663-9. PubMed PMID: 21869714.
- 95 Samuels Y, Wang Z, Bardelli A, et al. High frequency of mutations of the PIK3CA gene in human cancers. *Science*. 2004 Apr 23;304(5670):554. Epub 2004 Mar 11. PubMed PMID: 15016963.
- 96 Sanchez R, Ayala R, Martinez-Lopez J. Minimal residual disease monitoring with next-generation sequencing methodologies in hematological malignancies. *Int J Mol Sci*. 2019 Jun 10;20(11). PubMed PMID: 31185671.
- 97 Sartori DA, Chan DW. Biomarkers in prostate cancer: what's new? *Curr Opin Oncol*. 2014 May;26(3):259-64. PubMed PMID: 24626128.
- 98 Scher HI, Graf RP, Schreiber NA, et al. Assessment of the validity of nuclear-localized androgen receptor splice variant 7 in circulating tumor cells as a predictive biomarker for castration-resistant prostate cancer. *JAMA Oncol*. 2018 Sep 1;4(9):1179-1186. doi: 10.1001/jamaoncol.2018.1621. PubMed PMID: 29955787.
- 99 Senkus E, Kyriakides S, Ohno S, et al. Primary breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2015; 26(Suppl 5): v8-v30. PubMed PMID: 26314782.
- 100 Sestak I, Buus R, Cuzick J, et al. Comparison of the performance of 6 prognostic signatures for estrogen receptor-positive breast cancer: A secondary analysis of a randomized clinical trial. *JAMA Oncol*. 2018 Apr 1;4(4):545-553. PMID: 29450494.
- 101 Soekhojo CY, de Mel S, Ooi M, Yan B, Chng WJ. Potential Clinical Application of Genomics in Multiple Myeloma. *Int J Mol Sci*. 2018 Jun 10;19(6). pii: E1721. PubMed PMID: 29890777.
- 102 Song L, Li Y. SEPT9: A specific circulating biomarker for colorectal cancer. *Adv Clin Chem*. 2015; 72:171-204. Epub 2015 Aug 29. PubMed PMID: 26471083.
- 103 Sparano JA, Gray RJ, Makower DF, et al. Prospective validation of a 21-gene expression assay in breast cancer. *N Engl J Med* 2015; 373:2005-2014. PMID: 26412349.
- 104 Sparano JA, Gray RJ, Makower DF, et al. Adjuvant Chemotherapy Guided by a 21-Gene Expression Assay in Breast Cancer. *N Engl J Med*. 2018 Jul 12;379(2):111-121. PMID: 29860917.
- 105 Sparano JA, Gray RJ, Ravdin PM, et al. Clinical and genomic risk to guide the use of adjuvant therapy for breast cancer. *N Engl J Med*. 2019 Jun 20;380(25):2395-2405. PubMed PMID: 31157962.
- 106 Spivak JL. Myeloproliferative neoplasms. *N Engl J Med*. 2017 Jun 1;376(22):2168-81. PubMed PMID: 28564565.
- 107 Steward DL, Carty SE, Sippel RS, et al. Performance of a multigene genomic classifier in thyroid nodules with indeterminate cytology: A prospective blinded multicenter study. *JAMA Oncol*. 2018 Nov 8. doi: 10.1001/jamaoncol.2018.4616. [Epub ahead of print] PubMed PMID: 30419129.
- 108 Tefferi A. Primary myelofibrosis: 2017 update on diagnosis, risk-stratification, and management. *Am J Hematol*. 2016 Dec;91(12):1262-71. PubMed PMID: 27870387.
- 109 Umemura S, Tsuchihara K, Goto K. Genomic profiling of small-cell lung cancer: the era of targeted therapies. *Jpn J Clin Oncol*. 2015 Jun;45(6):513-9. Epub 2015 Feb 10. PubMed PMID: 25670763.
- 110 US Food & Drug Administration. PCA3 Assay. Summary of Safety and Effectiveness. Available at: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P100033>. Accessed on 9.23.2020.
- 111 Van Neste L, Partin AW, Stewart GD, Risk score predicts high-grade prostate cancer in DNA-methylation positive, histopathologically negative biopsies. *Prostate*. 2016 Sep;76(12):1078-87. PubMed PMID: 27121847.
- 112 Vannucchi AM, Barbui T, Cervantes F, et al.; ESMO Guidelines Committee. Philadelphia chromosome-negative chronic myeloproliferative neoplasms: ESMO Clinical Practice Guidelines for diagnosis, treatment, and follow-up. *Ann Oncol*. 2015 Sep;26 Suppl 5: v85-99. PubMed PMID: 26242182.
- 113 Varga Z, Sinn P, Seidman AD. Summary of head-to-head comparisons of patient risk classifications by the 21-gene Recurrence Score® (RS) assay and other genomic assays for early breast cancer. *Int J Cancer*. 2019 Jan 17. PubMed PMID: 30653259.
- 114 Vatandoost N, Ghanbari J, Mojaver M, et al. Early detection of colorectal cancer: from conventional methods to novel biomarkers. *J Cancer Res Clin Oncol*. 2016 Feb;142(2):341-51. Epub 2015 Feb 17. PubMed PMID: 25687380.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

- 115 Wong KS, Angell TE, Strickland KC, et al. Noninvasive follicular variant of papillary thyroid carcinoma and the Afirma gene-expression classifier. *Thyroid*. 2016 Jul; 26(7):911-5. PubMed PMID: 27219469.
- 116 Wong WJ, Pozdnyakova O. Myeloproliferative neoplasms: Diagnostic workup of the cythemic patient. *Int J Lab Hematol*. 2019 May;41 Suppl 1:142-150. PubMed PMID: 31069979.
- 117 Wood B, Wu D, Crossley B, Dai Y, Williamson D, Gawad C, Borowitz MJ, Devidas M, Maloney KW, Larsen E, Winick N, Raetz E, Carroll WL, Hunger SP, Loh ML, Robins H, Kirsch I. Measurable residual disease detection by high-throughput sequencing improves risk stratification for pediatric B-ALL. *Blood*. 2018 Mar 22; 131(12):1350-1359. PubMed PMID: 29284596.
- 118 Wu YL, Tsuboi M, He J, John T, Grohe C, Majem M, Goldman JW, Laktionov K, Kim SW, Kato T, Vu HV, Lu S, Lee KY, Akewanlop C, Yu CJ, de Marinis F, Bonanno L, Domine M, Shepherd FA, Zeng L, Hodge R, Atasoy A, Rukazenkov Y, Herbst RS; ADAURA Investigators. Osimertinib in Resected EGFR-Mutated Non-Small-Cell Lung Cancer. *N Engl J Med*. 2020 Oct 29;383(18):1711-1723. PMID: 32955177.
- 119 Zaman A, Bivona TG. Emerging application of genomics-guided therapeutics in personalized lung cancer treatment. *Ann Transl Med*. 2018 May;6(9):160. PubMed PMID: 29911108.
- 120 Zhang L, Zeng J, Zeng Z, et al. MGMT in colorectal cancer: a promising component of personalized treatment. *Tumour Biol*. 2016 Aug;37(8):11443-56. Epub 2016 Mar 22. PubMed PMID: 27006309.

Revision History

Medical Advisory Board Review:

- v2.2021 03/12/2021: Approved
- v1.2021 11/13/2020: Approved
- v4.2020 12/29/2020: Approved
- v3.2020 11/13/2020: Approved
- v2.2020 05/08/2020: Reviewed
- v1.2020 11/04/2019: Approved
- v3.2019 09/10/2019: Approved
- v2.2019 05/23/2019: Approved
- v1.2019 11/07/2018: Reviewed
- v1.2018 03/31/2018: Reviewed

Clinical Steering Committee Review:

- v2.2021 02/22/2021: Approved
- v1.2021 10/13/2020: Approved
- v4.2020 12/29/2020: Approved
- v3.2020 10/13/2020: Approved
- v2.2020 04/06/2020: Approved
- v1.2020 10/11/2019: Approved

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

v3.2019 09/09/2019: Approved

v2.2019 04/03/2019: Approved

v1.2019 10/03/2018: Approved

v1.2018 02/28/2018: Approved

v5.2017 11/01/2017: Approved

v4.2017 09/20/2017: Approved

v3.2017 08/09/2017: Approved

v2.2017 05/03/2017: Approved

v1.2017 01/25/2017: Approved

Revisions:

Version	Date	Editor	Description
v2.2021 GEN04-0921.1	2/15/2021	Heather Dorsey, MS, CGC	Semi-annual review. Formatting changes were made to Table 1 and T-Cell antigen receptor (TCR) was added for T-Cell Lymphoma (peripheral). Test name was corrected for ExoDx. CPT codes, professional society guidelines, background and references were updated.
v1.2021	9/11/2020	Heather Dorsey, MS, CGC	Semi-annual review. Criteria was added for cholangiocarcinoma and neuroblastoma testing. Prostate cancer and tumor agnostic criteria was revised. Breast cancer GEC, MRD testing and targeted testing for NTRK fusions criteria were updated. CPT codes, professional society guidelines, background and references were updated.
v4.2020	12/29/2020	Heather Dorsey, MS, CGC	Interim Update: Coverage criteria was expanded for OncotypeDX Breast Recurrence Score test. Coverage for EGFR in NSCLC (Stage IB-IIIa) was added.
v3.2020	10/9/2020	Heather Dorsey, MS, CGC	Interim Update: Coverage criteria was added for liquid biopsy testing in patients with metastatic castrate-resistant prostate cancer, metastatic breast cancer (updated), ovarian cancer and

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Version	Date	Editor	Description
			metastatic NSCLC. General coverage criteria for multi-gene panels was updated to clarify coverage for tests designated as FDA companion diagnostics. CPT codes, background and references were updated.
v2.2020	03/13/2020	Heather Dorsey, MS, CGC	General coverage criteria for somatic multi-gene panels was updated to include criteria for an FDA companion diagnostic. Criteria was added for CMA testing for multiple myeloma. Targeted multi-gene panels were added for metastatic castration-resistant prostate cancer. Gene list was updated for B-Cell Lymphoma and RET fusions were added for thyroid cancer. Gene expression classifier testing criteria for breast cancer was expanded. Prostate Cancer (symptomatic cancer screening) was clarified. Updated CPT codes, professional society guidelines, background and references.
v1.2020	10/02/2019	Heather Dorsey, MS, CGC	Clarification of cell free testing. Reformatted coverage criteria. Coverage criteria expansion for MPN to allow testing for JAK2, CALR, and MPL as well as criteria for targeted somatic testing of PIK3CA. Updated CPT codes, background, professional society guidelines and references.
	2/5/2020	Carrie Langbo, MS, CGC	NCCN Guidelines® were accessed for inclusion of the most recent published version. Minor revisions to text were incorporated based on updated Guidelines but did not impact coverage criteria/necessitate MAB/CSC review.
v3.2019	9/09/2019	Heather Dorsey, MS, CGC	Interim update. Minimal Residual Disease (MRD) testing criteria was added and coverage criteria for NTRK fusion testing was expanded to cover approved FDA medications. CPT codes, background, professional society guidelines and references were updated.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Version	Date	Editor	Description
v2.2019	4/03/2019	Emily Higuchi, MS, CGC	Semi-annual review. Revised umbrella coverage criteria section. Added NTRK fusion criteria. Revised Oncotype DX®, Prosigna PAM50™ and MammaPrint® criteria. Added Endopredict criteria. Updated background, professional society/NCCN® guidelines and references. Renumbered to v2.2019.
	7/25/2019	Carrie Langbo, MS, CGC	NCCN Guidelines® were accessed for inclusion of the most recent published version. Minor revisions to text were incorporated based on updated Guidelines but did not impact coverage criteria/necessitate MAB/CSC review.
v1.2019	03/04/2019	Gwen Fraley, MS, CGC	Urgent Interim review. Expand coverage of ThyroSeq3.0 for indeterminate thyroid nodules and revision to reflect current testing platforms.
v1.2019	11/01/2018	Ashley Allenby, MS, CGC	Semi-annual review. Removed NCCN® 2B criteria recommendation from general medical necessity criteria. Added criteria for ThyroSeq3.0. Updated background, professional society/NCCN Guidelines® and references. Renumbered to 2019. Reformatted CPT code list. PMID added.
v1.2018	03/31/2018	Gwen Fraley, MS, CGC	Semi-annual review. Added disclaimer sentence to scope section. Added uveal melanoma to list of tumor types for somatic genetic testing. Added exclusion criteria for prostate cancer tumor testing. Revised MammaPrint® criteria. Updated background, professional society/NCCN Guidelines and references. Renumbered to 2018. Submitted to CSC for approval.
v5.2017	11/01/2017	Gwen Fraley, MS, CGC	Revised criteria for indeterminate thyroid nodules. Updated background and references. Renumbered to v5.2017 and submitted to CSC for approval.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Version	Date	Editor	Description
v4.2017	09/18/17	Megan Czarniecki, MS, CGC	Removed specific criteria for lung cancer. Formatting changes: converted references to NLM style. Incorporated “methodological considerations” to appropriate use criteria and background. Renumbered to v4.2017 and submitted to CSC for approval.
v3.2017	08/09/2017	Gwen Fraley, MS, CGC	Changed nomenclature of “occult primary” to “cancer of unknown primary/occult neoplasm”. Changed stance on MammaPrint® to allow for coverage when criteria met. Removed separate lung cancer criteria and referred to NCCN. Updated references. Added additional codes to Coding Considerations.
v2.2017	06/30/2017	Denise Jones, MS, CGC	Quarterly review. No criteria changes. Updated references.
v2.2017	04/25/2017	Cheryl Thomas, MS, CGC	Quarterly review. Added changes to indeterminate thyroid nodules (removed Hurthle cell from indication per NCCN update). Added PD-L1 to NSCLC molecular targets. Updated references.
v1.2017	01/23/2017	Gwen Fraley, MS, CGC	Quarterly review. Updated MPN criteria. Edited EGFR criteria regarding erlotinib. Updated references. Renumbered to 2017.
v4.2016	09/29/2016	Jenna McLosky, MS, CGC	Updated background regarding occult primaries. Updated references.
v3.2016	06/30/2016	Jenna McLosky, MS, CGC	Added EGFR Cobas cell-free test for NSCLC. Updated references.
v2.2016	04/04/2016	Jenna McLosky, MS, CGC	Updated and reviewed prostate cancer screening criteria. Updated references.

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.

Version	Date	Editor	Description
v1.2016	03/18/2016	Jenna McLosky, MS, CGC	Updated and revised stance on breast cancer prognosis assays (Prosigna). Updated references.
v1.2015	09/24/2015	Jenna McLosky, MS, CGC	Original version

Original Effective Date: 09/24/2015

Primary Author: Jenna McLosky, MS, CGC

ARCHIVED

PROPRIETARY

Guidelines developed by, and used with permission from, Informed Medical Decisions, Inc. © 2015-2021 Informed Medical Decisions, Inc. All Rights Reserved.